

A monographic study of the genus *Sparganium* (Sparganiaceae). Part 1, Subgenus *Xanthosparganium* Holmberg

Autor(en): **Cook, Christopher D.K. / Nicholls, Marc S.**

Objekttyp: **Article**

Zeitschrift: **Botanica Helvetica**

Band(Jahr): **96(1986)**

Heft 2

Erstellt am: **Apr 28, 2013**

Persistenter Link: <http://dx.doi.org/10.5169/seals-67202>

Nutzungsbedingungen

Mit dem Zugriff auf den vorliegenden Inhalt gelten die Nutzungsbedingungen als akzeptiert. Die angebotenen Dokumente stehen für nicht-kommerzielle Zwecke in Lehre, Forschung und für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und unter deren Einhaltung weitergegeben werden. Die Speicherung von Teilen des elektronischen Angebots auf anderen Servern ist nur mit vorheriger schriftlicher Genehmigung des Konsortiums der Schweizer Hochschulbibliotheken möglich. Die Rechte für diese und andere Nutzungsarten der Inhalte liegen beim Herausgeber bzw. beim Verlag.

SEALS

Ein Dienst des Konsortiums der Schweizer Hochschulbibliotheken
c/o ETH-Bibliothek, Rämistrasse 101, 8092 Zürich, Schweiz
retro@seals.ch
<http://retro.seals.ch>

A monographic study of the genus *Sparganium* (Sparganiaceae).

Part 1. Subgenus *Xanthosparganium* Holmberg

Christopher D. K. Cook and Marc S. Nicholls

Institut für Systematische Botanik der Universität Zürich, Zollikerstr. 107, CH-8008 Zürich,
Switzerland

Manuscript accepted September 9, 1986.

Abstract

Cook, C. D. K. & Nicholls, M. S. 1986. A monographic study of the genus *Sparganium*.
Part 1. Subgenus *Xanthosparganium* Holmberg. *Bot. Helv.*, 96: 213–267.

A taxonomic revision of the genus *Sparganium* (Sparganiaceae) is presented with a key and full descriptions of each species including diagnoses, synonyms with typifications, distribution maps, variation and illustrations also containing information on fossils, anatomy, morphology, chromosomes, floral biology, dispersal, ecology, parasites, animal feeders, applied aspects and hybrids. Fourteen species and six subspecies are recognised. In part 1, subgenus *Xanthosparganium*, seven species are presented. One new name is used: *Sparganium emersum* subsp. *acaule* (Beeby ex Macoun) C. D. K. Cook & M. S. Nicholls.

Introduction

The only attempt at a worldwide revision of *Sparganium* was written by Graebner (1900). It was based almost entirely on the herbarium material in Berlin, most of which was destroyed in the second world war. Graebner worked in the Englerian fashion and recognised numerous taxa at varying ranks: 3 sections, 16 living and 16 fossil species, 4 subspecies, 5 proles, 28 varieties, 3 subvarieties and 4 hybrids. He described 12 new taxa and made 16 new nomenclatural combinations. Like Linneus before him, Graebner based his major taxa on vegetative characters. As virtually all species can grow submerged, floating and emergent many of Graebner's taxa represent no more than phenotypic states of a single taxon. Graebner also based some taxa on discordant elements, an additional source of confusion. This revision has simply not stood the test of time.

Wladislaw Rothert (b. 1863) formerly of the University of Kasan, later at Charkow, Odessa, Riga, Bogor, Warsaw and Krakow was preparing a monograph of *Sparganium*. He died in St. Petersburg (Leningrad) in 1916 before his monograph was published. We have not been able to trace his manuscript and it seems likely that it was destroyed during the upheavals in Russia at the time of Rothert's death. Two short accounts were published, one in 1910 on the species in Russia and the other in 1913 in Fedtschenko,

B. A., Flora Aziatskoi Rossii, 1: 17–37 St. Petersburg. Rothert made numerous notes on herbarium sheets in many different herbaria; we have often found these notes valuable for our studies. Rothert's herbarium is deposited in the Polska Akademia Nauk in Krakow; unfortunately, we have not been able to see this extensive collection but it apparently contains no holotypes and only three specimens collected by Rothert himself. C. D. K. Cook started studying *Sparganium* in 1956 and wrote some minor accounts in 1961, 1962 and 1963. After a quarter of a century's dormancy the genus has been, once more, actively studied.

Materials and methods

Our work is based mainly on herbarium and library studies with some field studies and experiments on living plants. Except for types and some otherwise important plants we have not cited herbarium sheets. We can give information about the material we have examined on request. We have also labelled herbarium specimens we have examined. Material from the following herbaria has been seen by us: State Herbarium of South Australia, Adelaide, Australia (AD); Botanischer Garten und Botanisches Museum, Berlin-Dahlem, W. Germany (B); Queensland Herbarium, Indooroopilly, Australia (BRI); Herbarium Australiense, Canberra City, Australia (CANB); Royal Botanic Garden, Edinburgh, UK (E); Conservatoire et Jardin Botaniques, Genève, Switzerland (G); Gray Herbarium of Harvard University, Cambridge, USA (GH); Royal Botanic Gardens, Kew, UK (K); Landesmuseum für Kärnten, Klagenfurt, Austria (KL); Kyoto University, Kyoto, Japan (KYO); Botanical Museum, Lund, Sweden (LD); Botanische Staatssammlung, München, W. Germany (M); Makino Herbarium, Metropolitan University, Tokyo (MAK); National Herbarium of Victoria, Melbourne, Australia (MEL); Lomonosov State University, Moscow, USSR (MW); New England Botanical Club Herbarium, Cambridge, USA (NEBC); National Herbarium of New South Wales, Royal Botanic Gardens, Sydney (NSW); Naturhistoriskariksmuseet, Stockholm, Sweden (S); University of Tokyo, Hongo, Tokyo, Japan (TI); University of Uppsala, Uppsala, Sweden (UPS); Institut für Systematische Botanik der Universität, Zürich, Switzerland (Z); Eidg. Technische Hochschule, Zürich, Switzerland (ZT).

Photographs of types and other important specimens were generously supplied by: the Komarov Botanical Institute of the Academy of Sciences, Leningrad, USSR (LE); New York Botanical Garden, New York, USA (NY).

Acknowledgements

We would like to thank the Fonds National Suisse de la Recherche Scientifique (Nr. 3.390-0.83) for a post-doctoral fellowship for Marc Nicholls. We would also like to thank the following for material or specialized knowledge: Helen Aston, Melbourne, Australia; Adolf Czeska, Victoria, B.C., Canada; Urs Eggli, Zürich, Switzerland; V. I. Grubov, Leningrad, USSR; Heino Heine, Paris, France; Patricia Holmgren, New York, USA; A. Jasiewicz, Krakow, Poland; A. N. Krasnova, A.N., Borok, USSR; Per Lassen, Lund, Sweden; Lian Guang-Hua, Nanjing, China; Li Hen, Kunming, China; John Mc Neill, Ottawa, Canada; Roland Moberg, Uppsala, Sweden; Bertil Norstenstam, Stockholm, Sweden.

We would also like to thank the curators of the herbaria listed above for lending us their material.

Special terminology

In previous publications on *Sparganium* some terms have been widely used and have proved helpful although some are technically incorrect. The following specialized terms are used in this account:

Approximate – Referring to heads which overlap each other.

Axillary – Arising in the axis of a bract, exactly where the bract departs from the main stem, Fig. 1 (see also *Supra-axillary*).

Beak – The persistent part of the style.

Bract – A leaf subtending an inflorescence branch or head; it may be green and leaf-like or brownish and scale-like (Fig. 1).

Branch – A lateral stem bearing more than one head, Fig. 1 (see also *Peduncle*).

Endocarp – The innermost differentiated layer of pericarp, in other works often incorrectly called seed. Measurements made without the stalk.

Fruit – Measurements and shape descriptions made excluding the beak and the pedicel.

Head – An almost spherical, crowded group of sessile or subsessile flowers sharing a common base, in other works often called capitulum.

Inflorescence – The part of the plant bearing heads (see also *Head*).

Peduncle – The stalk bearing a single head (Fig. 1). The usage is technically incorrect (see p. 221) because the heads are compound inflorescences.




Fig. 1. Diagrammatic representation of *Sparganium* inflorescences to illustrate special terminology:
 1. leaf; 2. lowest bract; 3. axillary branch; 4. axillary peduncle; 5. axillary head; 6. supra-axillary peduncle; 7. supra-axillary head; 8. supra-axillary head concoulescent above the next node; 9. approximate heads.

Perianth segments – The scale-like segments surrounding each flower. In other works sometimes called tepals; no distinction is made between bracts, bractioles, sepals or petals, see also p. 221.

Remote – Referring to heads which do not touch each other.

Supra-axillary – Arising on the main stem above the point where the bract departs from the main stem; branch, peduncle or head fused to main stem (concaulescent). Occasionally concaulescent up to or even above the next node, Fig. 1.

Classification

Sparganiaceae F. Rudolphi, *Systema Orbis Vegetabilium*, 27. October 1830.

One genus: *Sparganium*.

Following exhaustive investigations into the structure of the flower and inflorescence of *Sparganium* and *Typha*, Müller-Doblies, D. (1970) was convinced that *Sparganium* is so closely related to *Typha* that both should be united into the family *Typhaceae* (A. L. de Jussieu, *Genera Plantarum*, 25 October 1789).

We acknowledge that *Sparganium* and *Typha* share numerous features and that there is evidence of patristic relationship. However, they are clearly distinct from each other and have been so since, at least, the Oligocene (we rather distrust the records for both *Sparganium* and *Typha* from the Cretaceous). We therefore see no purpose in uniting both families; in fact, to do so will quite unnecessarily increase nomenclatural instability as nearly all floristic works in the last 150 years have recognized the two families as distinct.

Sparganium L. *Species Plantarum*, 971. 1 May 1753.

Type: Rydberg (N. Amer. Flora, 17: 5. 30 June 1909) designated *S. erectum* the type of the genus. This is unfortunate as *S. erectum* is based on a somewhat incomplete illustration in l'Obel, *Plantarum Seu Stirpium Historia*, 41. 1576 (see Cook 1985). We would prefer to base the genus *Sparganium* on *S. natans* which is, at least, based on an adequate specimen (sheet 1095/2, in Linn, see p. 234 and Cook 1985).

The following species were incorrectly described in the genus *Sparganium*; details are given by Cook (1985):

- S. tenuifolia* [sic] Poiret, *Voyage en Barbarie*, 2: 253. 1788, has been transferred to the genus *Carex* (under *Carex halleriana* Asso), see op. cit., p. 317.
- S. pubescens* Poiret, op. cit., has been transferred to the genus *Fuirena* (under *F. pubescens* (Poiret) Knuth).
- S. trifidum* Poiret in Poiret & Lamarck, *Encyclopédie Méthodique, Botanique*, Suppl. 4, 723. 1816, has been transferred to the genus *Dulichium* (under *D. arundinaceum* (L.) Britton).

Family and generic description

Aquatic or amphibious, monoecious, glabrous, perennial herbs.

Vessels with scalariform perforation plates; sieve tube plastids P-type with numerous cuneate protein crystalloids; oxalate raphides frequent; silica bodies absent.




Fig. 2. *Sparganium*: a, perianth segments (left: *S. erectum* subsp. *neglectum*; right: *S. emersum* scale bar: 1 mm); b, adaxial surface of leaf base of *S. erectum* subsp. *neglectum* showing glands, (scale bar: 1 cm); c, pollen grain of *S. erectum* subsp. *neglectum* (Scale bar: 5 µm); d, surface of pollen grain of *S. fluctuans* (scale bar: 1 µm).

Roots adventitious, fibrous, often inflated, simple or branched; root hairs not arising from specialised cells.

Stems stoloniferous to rhizomatous or cormous below, erect or floating and usually becoming emergent above, sympodial with reiterative monopodia, each monopodial part living for not more than 2 years.

Leaves distichous, exstipulate, sheathing below (sheath often tubular at base), not differentiated into petiole with blade, scale-like on underground parts otherwise linear; scale-like leaves brownish, translucent, thin and mostly remote; linear leaves submerged, floating or emergent, mostly basal and equitant or cauline and remote, more or less flattened to V-shaped or triangular in transverse section, sometimes carinate or keeled, with many longitudinal veins connected by transverse diaphragms; air cham-

bers in 1 or usually more layers; stomata paracytic, the surrounding cells with sometimes intersecting oblique divisions; margins entire; apex rounded or apiculate.

Flowers unisexual, arranged tightly in 2 or 3 compact racemes in globose heads; heads usually unisexual, sessile or pedunculate, axillary or supra-axillary (concaulescent), or occasionally concaulescent up to or beyond the next node (Fig. 1), arranged in simple or branched racemes with female heads proximal to the males or occasionally the lowest branch bearing male heads only; male heads usually remote below becoming compacted above, caducous but naked axis usually persisting; inflorescence bracts leaf-like below, becoming reduced and scale-like in the middle and eventually absent above.

Floral bract usually indistinguishable from perianth segments; bracteoles absent.

Perianth segments radiate, (1-)3-4(-6), dry, scale-like, usually widened above (Fig. 2 a), often irregular in shape and arrangement, sometimes superposed and sometimes united below.

Female flowers without staminodia, without nectaries; ovary superior of 1-3 (in fossil species to 7, Fig. 4a) united carpels; style 1, beak-like, often persisting in fruit or reduced; stigmas 1-3 (or perhaps more in fossil species), papillose, dry; ovules solitary, ventrally anatropous, pendulous, bitegmic with the outer integument longer than the inner, crassinucellar; megagametophyte development Polygonum-type; embryogenesis Onograd-type endosperm Helobial; embryo linear, straight and large, without chlorophyll.

Male flowers tightly packed in heads and difficult to recognise as individual flowers; stamens 1-8, often superposed; filaments mostly free, elongate, falling after anthesis; connectives somewhat widened above; anthers basifix, oblong, 4-microsporangiate, dehiscing neutrally through longitudinal slits; endothecium with spiral thickening; tapetum 8-nucleate, at first amoeboidal, later pseudo-periplasmoidal; microsporogenesis successive; pollen single, monoporate, ulcerate, spheric to ellipsoidal (Fig. 2c), the surface reticulate (Fig. 2d), 2-cellular when released; pistillodia absent; nectaries absent.

Fruits sessile to shortly stalked, drupe-like, crowded in a burr-like head but released singly; mesocarp spongy; endocarp hard, with longitudinal ridges or smooth, with distinct micropylar plug.

Seeds with a membranous testa and starchy endosperm surrounding the embryo.

Distribution

The genus *Sparganium* is found almost throughout the temperate and arctic regions of the northern Hemisphere. In the Tropics there are isolated occurrences in the mountains of central Sumatra and New Guinea. In the southern temperate regions it occurs in southeastern Australia and New Zealand (see Fig. 3). Most species are found in the northern boreal zone and the regions of greatest species diversity are in North America and eastern Asia. There are no local endemics and nearly all species have large more or less continuous distribution patterns; only *S. glomeratum* (see p. 242) and *S. subglobosum* show major disjunctions. *Sparganium* seems to be absent from most of the Himalayas, being recorded only in the extreme west and extreme east. All fossil finds are within the range occupied by *Sparganium* today. Only one species seems to have become established outside its native area in recent times; it is *S. erectum* which is perhaps introduced in Australia.




Fig. 3. World-wide distribution of *Sparganium*. The numbers represent the number of species in that region.

Fossils

Records of pollen and fruits have been reported from the Cretaceous. These records are based on very doubtful evidence although Mai (1984) is inclined to accept the report of Velenovský & Viníklář (1926) of supposed fruiting heads of *Sparganium* from Czechoslovakia. The earliest reliable record of *Sparganium* in an imprint of an inflorescence from the Middle Eocene of Wyoming named *S. antiquum* (Newberry) Berry. In his paper Berry (1924) draws a reconstruction which we feel must be wrong, the lower portion based on *Pontederites hesperia* Knowlton should not have been combined with the inflorescence which is probably *Sparganium*. We have not, however, examined the original fossil material. Without qualification Muller (1984) cites the Paleocene for the first *Sparganium* fruits and for Typhales pollen.

By the Oligocene there is fossil material from the Old World that is convincingly attributable to the genus *Sparganium*, both pollen (Abuziarova 1970) and macrorests. A superficial search through the literature has revealed that there are at least 44 fossil species of *Sparganium* described. At least 34 of them are certainly *Sparganium* but we are not convinced they are all distinct from each other or that they are all distinct from recent species.

Among the fossil species both subgenera (*Sparganium* and *Xanthosparganium*) are already distinct by the Oligocene. Most of the Oligocene species are clearly distinct from the species of today. The endocarps although shorter than some recent species were often much wider with numerous locules (*S. balticum* Dorofeev (Fig. 4a), 4–7, locules; *S. multiloculare* Reid & Chandler, 2–5; *S. sobolevii* Dorofeev, 2–4, but surface smooth). Another species *S. costatum* Dorofeev from Oligocene beds near Tomsk had a




Fig. 4. Drawings of *Sparganium* endocarps: a, *S. balticum*; b, *S. costatum*; c, *S. erectum* subsp. *neglectum*; d, *S. emersum* (scale bar: 1 mm).

winged endocarp quite unlike any recent species (Fig. 4b). By the Miocene the endocarps of the majority of the described fossil species are very difficult to distinguish from recent species and even Dorofeev (1979) who has himself described no less than 33 fossil species recognises *S. erectum* (Fig. 4c) as going back to the Miocene.

Anatomy and morphology

a) Vegetative parts

The general anatomy and morphology of the vegetative parts of *Sparganium* are well-summarized by Loew in Kirchner et al. (1908) and by Meyer in Solereder & Meyer (1933); the essential features of diagnostic value have been incorporated into the formal descriptions.

The growth is essentially sympodial, each horizontal axis (rhizome or stolon depending on the species) is a monopodium which bends upright at the tip where it becomes thickened and develops into a hard rather woody corm. After the first winter each corm usually develops several axillary monopodia and then dies with or without developing an inflorescence. Photosynthetic leaves develop only from corms; the rhizomes or stolons bear non-photosynthetic, scale-like leaves. The leaves are distichous which has the consequence that the lateral monopodia tend to leave the mother corm at angles approaching 180°.

The lateral monopodia grow horizontally through their subtending leaves, they do not appear to be dorso-ventrally organised and do not, under normal conditions, branch. If damaged these horizontal axes are capable of further growth through the development of buds in the axils of the scale leaves. Stems and roots are ventilated internally by a system of large chambers bridged by porous diaphragms.

Germination takes place submerged in water. The juvenile leaves of all species are aquatic and cannot tolerate emergence. Depending upon the species and local ecological conditions the leaves may remain submerged or, later in ontogeny, become floating or emergent. During the juvenile phase it is not possible to distinguish the species. The submerged leaves have a relatively simple system of longitudinal compartments with porous transverse diaphragms. The compartments between the diaphragms are partly schizogenous and partly lysigenous in origin.

The architecture of *Sparganium* leaves has been described by Kaul (1972, 1973, 1976). In a large and erect species such as *S. eurycarpum* the emergent leaves are deltate in transverse section with palisade tissue on all three sides. Within the leaf every fourth diaphragm has a vascular bundle and is heavily photosynthetic, the other diaphragms are translucent and consist of thin stellate cells (actinochyma). Floating leaves are isobilateral with adaxial palisade only and there are even more thin, essentially non-photosynthetic diaphragms than thick photosynthetic ones.

If all diaphragms were thick and heavily photosynthetic and had small intercellular spaces, the amount of light reaching the interior would be diminished and as Kaul (1972) suggests this pattern of alternating translucent and photosynthetic diaphragms allows the linear leaves of *Sparganium* with a relatively small surface area to utilise more light than leaves without such internal structures.

On the adaxial surface of the upper portion of the sheathing part of the leaf there are curious longitudinal rows of emergent glandular cells (Fig. 2 b). These glands are brownish and are sometimes visible to the naked eye. In *Typha* similar looking glands secrete slime, this does not appear to be the case in *Sparganium*. The statement by Graebner (1900) that these glands are lacking in *Sparganium* is false.

b) *Floral parts*

The morphology of the floral parts of *Sparganium* has been exhaustively described by Müller-Doblies, Ute (1969), Müller-Doblies, Dietrich (1970) and Müller-Doblies, Ute & Müller-Doblies, Dietrich (1977). The primary inflorescence is a simple or branched raceme, the female and lateral male heads are twice racemose but the terminal male heads are three-times racemose or consist of several virtually united lateral heads. The structure of the heads can be followed only by examination of the total ontogeny.

The female flowers have as a rule one bract and (2-)3-4(-5) perianth segments. There is no fundamental difference between the bract and the perianth segments (Fig. 2 a); differences such as size, innervation and ontogenetic succession may be reduced and in extreme cases totally de-differentiated. The male flower is essentially like the female but has 1 to 8 stamens and lacks traces of pistillodium.

The heads are usually unisexual but some female flowers in essentially male heads are not uncommon, they are, however, very rarely fertile. Heads partly male and partly female are uncommon. Intersex flowers are found from time to time but the female component is usually sterile. The plants are usually bisexual but in most species pure male plants have been found.

Although morphologically incorrect, the term "branch" is used in this and all other known accounts for axillary shoots bearing more than one head and the term "peduncle" for axillary shoots terminating in solitary heads. The "peduncles" are strictly speaking lateral branches because they are branched below the inflorescence, each head is, strictly speaking, made up of two or three compact racemes (dicephalus or tricephalus). Genuine terminal heads were not found by Müller-Doblies, Ute (1969).

Chromosomes

The published chromosome numbers are presented on Table 1; because of the high uniformity no efforts have been made to check voucher specimens and we have not made any chromosome investigations ourselves. About 65 plants have been counted

Table 1. *Sparganium* chromosome numbers.

Name (in publication)	Number	Locality	Voucher	III.	Reference
S. angustifolium					
(<i>angustifolium</i>)	2n=30	None given, presumably Scandinavia	?	-	Löve & Löve 1942
(<i>angustifolium</i>)	2n=30	Iceland	?	-	Löve & Löve 1956
(<i>affine</i>)	2n=30	None given, presumably Poland	?	?	Bijok & Mlynek 1965
(<i>angustifolium</i>)	2n=30	Canada: Manitoba, Machbride Lake, s.d., <i>J. C. Ritchie 1075</i>	WIN	-	Löve & Ritchie 1966
(<i>multipedunculatum</i>)	2n=30	Canada: Manitoba, New Iceland, Vannes, s.d., <i>Löve & Löve 5566</i>	WIN	-	Löve & Löve 1975
(<i>angustifolium</i>)	2n=30	Czechoslovakia: Slovákovia, Západné Tatry, Rohačské plesá, s.d., Májovský & Murin s.n.	SLO	-	Murin & Májovský 1978b
(<i>angustifolium</i>)	2n=30	Canada: Manitoba, Brokenhead River, near Vivian, s.d., Löve & Löve 5272	WIN or COLO	-	Löve & Löve 1981
S. emersum					
(<i>simplex</i>)	2n=30	Germany: Schleswig-Holstein	?	?	Scheerer 1940
(<i>simplex</i>)	2n=30	None given, presumably Scandinavia	?	?	Hagerup 1941
(<i>simplex</i>)	2n=30	None given, presumably Scandinavia	-	-	Löve & Löve 1942
(<i>simplex</i>)	2n=30	USSR: Kamchatka, near Kozytresvk, 24 July 1959, <i>Sokolovskaya</i> s.n.	LECB	-	Sokolovskaya 1963
(<i>simplex</i>)	2n=30	Poland	?	?	Bijok & Mlynek 1965
(<i>simplex</i>)	2n=30	Canada: Queen Charlotte Islands, ? coll.	?	?	Taylor & Mulligan 1968
(<i>simplex</i>)	2n=30	USSR: Leningrad Distr., Lake Buoksa, s.d., <i>Sokolovskaya</i> s.n.	LECB	-	Sokolovskaya 1972
(<i>emersum</i>)	2n=30	Czechoslovakia: Západné Beskydy, Zázrivá, s.d.,	SLO	-	Váchová in Májovský 1976
S. erectum s.l.					
(<i>ramosum</i>)	2n=30	Germany: Schleswig-Holstein, an der Kronsbek bei Neudorf, s.d., ? coll.	?	+	Wulff 1938
(<i>ramosum</i>)	2n=30	None given, presumably Scandinavia	-	-	Löve & Löve 1942
(<i>ramosum</i>)	2n=30	None given, presumably Poland	?	?	Bijok & Mlynek 1965
(<i>erectum</i>)	2n=30	Denmark: Bestmose Isl. of Bornholm, s.d., <i>K. Larsen 6039-Bn19</i>	AAU	-	Larsen 1965
(<i>erectum</i>)	2n=30	Netherlands: Utrecht, Vuntus, nr. Loodrecht, s.d., <i>Gadella & Kliphuis</i> s.n.	U	-	Gadella & Kliphuis 1973

(<i>erectum</i>)	2n=30	Czechoslovakia: Jur pri Bratislave, Podunajská nižina, Šur, s.d., <i>V. Feráková s.n.</i>	SLO	-	Feráková in Májovský 1974
(<i>ramosum</i>)	2n=30	India: Kashmir, Nagin Lake, s.d., <i>Mehra & Pandita 73</i>	PAN	-	Mehra & Pandita 1979
subsp. <i>erectum</i> <i>(polyedrum)</i> <i>(erectum)</i>	2n=30	None given	Löve & Löve 1948	-	Löve & Löve 1948
	2n=30	Czechoslovakia: Bratislava, Podunajská nižina, s.d., <i>V. Feráková s.n.</i>	SLO	-	Murin in Mákovský 1978 a
subsp. <i>microcarpum</i> <i>(microcarpum)</i>	2n=30	None given	-	-	Löve & Löve 1948
subsp. <i>neglectum</i> <i>(neglectum)</i> <i>(neglectum)</i>	2n=30	None given	-	-	Kožuharov & Kuzmanov 1964
	2n=30	Bulgaria: Smoljan, "Big Lake", 13 Sept. 1963 ? coll.	?	-	
(<i>neglectum</i>) <i>(neglectum)</i>	2n=30	Poland: pow. Cieszyn, Debowiec, s.d., ? coll.	?	+	Bijok & Adamkiewicz 1971
	2n=30	Poland: pow. Cieszyn, stawy Ochaby, s.d., ? coll.	?	+	Bijok & Adamkiewicz 1971
(<i>neglectum</i>)	2n=30	Poland: pow. Milicz, stawy Milicz, s.d., ? coll.	?	+	Bijok & Adamkiewicz 1971
subsp. <i>stoloniferum</i> <i>(stoloniferum)</i> <i>(stoloniferum)</i>	2n=30	None given, presumably Japan	-	+	Harada 1949
	n=15 II	None given, presumably Japan	-	+	Harada 1949
S. <i>eurycarpum</i>	2n=30	Canada: Manitoba, Brokenhead River, Vivian, Löve & Löve s.n.	WINor COLO	-	Löve & Löve 1981
subsp. <i>coreanum</i> <i>(macrocarpum)</i> <i>(coreanum)</i>	2n=30	None given, presumably Japan	-	+	Harada 1942
	2n=30	None given, presumably Japan	-	+	Harada 1942
S. <i>fallax</i> <i>(yamatense)</i> <i>(yamatense)</i>	2n=30	None given, presumably Japan	-	+	Harada 1949
	n=15 II	None given, presumably Japan	-	+	Harada 1949
S. <i>glomeratum</i> <i>(glomeratum)</i> <i>(glomeratum)</i>	2n=30	None given, presumably Scandinavia	-	-	Löve & Löve 1942
	2n=30	None given, presumably Japan	-	+	Harada 1949
S. <i>gramineum</i> <i>(friesii)</i>	2n=30	None given, presumably Scandinavia	-	-	Löve & Löve 1942

Table 1 (continued)

Name (in publication)	Number	Locality	Voucher	III.	Reference
S. hyperboreum					
(hyperboreum)	2n=30	None given, presumably Iceland	—	—	Löve & Löve 1948
(hyperboreum)	2n=30	Greenland	?	?	Jörgen et al. 1958
(hyperboreum)	2n=30	Canada: Queen Charlotte Islands, s.d., ? coll.	?	?	Taylor & Mulligan 1968
(hyperboreum)	2n=30	USSR: Anadyrskoe, Nagorbe, R. Komarinasa, s.d., ? coll.	?	—	Zhukova & Tikhonova 1971
(hyperboreum)	2n=30	USSR: Rud Jagodnyj, s.d., ? coll.	?	—	Zhukova & Petrovsky 1976
(hyperboreum)	2n=30	USSR: northern Pekul'veyem River, s.d., ? coll.	?	—	Zhukova 1980
(hyperboreum)	2n=30	Canada: north-east	?	?	Gervais 1981
(hyperboreum)	2n=30	Canada: Manitoba, New Iceland, opposite Hekla Island, s.d., Löve & Löve 3329	WINor	—	Löve & Löve 1981
			COLO		
S. japonicum					
(japonicum)	2n=30	None given, presumably Japan	—	+	Harada 1949
(japonicum)	n=15 II	None given, presumably Japan	—	+	Harada 1949
S. natans					
(minimum)	2n=30	Germany: Schleswig-Holstein, Schwabstedter Moor, s.d., ? coll.	?	—	Wulff 1938
(minimum)	n=15	Iceland	G	—	Löve & Löve 1956
(minimum)	n=15	Canada: Queen Charlotte Islands, 9 August 1964, Calder & Taylor s.n.	?	—	Taylor & Mulligan 1968
(minimum)	2n=30	Poland: pow. Giżycko, Bagiennice, s.d., ? coll.	?	+	Bijok & Adamkiewicz 1971
(minimum)	2n=30	Poland: pow. Mrągowo, Lukniany, ? coll.	?	+	Bijok & Adamkiewicz 1971
(minimum)	2n=30	Czechoslovakia: Západné Beskydy, Zázriva, s.d., Králik & Váčhová s.n.	SLO	—	Váčhová in Májovský 1976
(minimum)	2n=30	Canada: Manitoba, Porcupine Mt., s.d., Löve & Löve 3280	COLO		Löve & Löve 1981
(minimum)	2n=30	Finland: Ab Nauvo, Seili Arohonka, 177208	TUR	—	Arohonka 1982
S. subglobosum					
(stenophyllum)	2n=30	Japan: Botanic Garden of Kyoto University, s.d., ? coll.	?	+	Harada 1949
(stenophyllum)	2n=45	Botanic Garden of Kyoto University, s.d., ? coll.	?	+	Harada 1949
(stenophyllum)	n=15 II	Japan: Prov. Kida of Sikoku, s.d., ? coll.	?	+	Harada 1949
(stenophyllum)	2n=30	Japan: Prov. Musasi, Inada-Inoborito s.d., ? coll.	?	+	Harada 1949
(stenophyllum)	n=15 II	Japan: Prov. Musasi, Inada-Inoborito, s.d., ? coll.	?	+	Harada 1949

and with one exception all have $2n=30$. The exception was a single sample of *S. subglobosum* from the botanic garden of Kyoto University, Japan; it had $2n=45$ but Harada (1949) mentioned that other plants of the same species from the same place had $2n=30$. Meiosis has been reported from six plants, all had 15 bivalents and no irregularities.

The chromosomes of *Sparganium* are (0.7–)0.9–1.5(–2.0) μm long and can be classified as relatively small. Some published illustrations show more or less equal-sized chromosomes while others show differentiation into shorter and longer. Most illustrations do not clearly show the centromere, however, Bijok & Adamkiewicz (1971) found about half the chromosomes to be metacentric while the other half was acrocentric with an arm ratio of about 1:3.

Floral biology

The flowers are congested into globose unisexual heads with males borne above females. Exceptionally, some essentially male heads have female or intersex flowers which rarely develop fruit; male flowers in essentially female heads are very rare. Occasionally part of the head is male and part female.

Within a head, anthesis is more or less simultaneous. The heads are clearly protogynous in spite of the fact that many early publications, including Graebner (1900) have reported *Sparganium* to be protandrous. In some of the large branched plants the lowest female head has frequently passed its receptive stage before the first male heads shed pollen. The lowest female heads sometimes bear less fruit than the others; this is particularly noticeable in *S. erectum*. No agamospermous or self-incompatibility mechanisms have been found.

Pollen transfer is by wind. *S. erectum* is occasionally visited by representatives of syrphid flies of the Melanostoma-Platycheirus group according to Leereveld (1984). Pollen is a principle source of food for these flies and analyses of the crop and gut contents showed that the syrphid *Pyrophaena granditarsa* may feed almost exclusively on the inflorescence of *S. erectum* in northwestern Holland. Pinkess (1980) noted that not only syrphids but that bumble bees and honey bees also collect *Sparganium* pollen. Pinkess (1980) and Leereveld (1984) claim that zoophilous pollination may sometimes result from this activity but have not tested it experimentally. We have observed insects collecting pollen on *S. erectum* but female heads are visited very rarely and then for very short times.

We have studied pollen flow in a stand of *S. erectum* in an artificial pool at the University of Zürich by emasculating plants at known distances from the nearest male inflorescence and recording the number of seeds set. Emasculated plants separated from the nearest male inflorescence by a distance of 1 m set less than 25% of the seeds set of intact plants, though even plants separated by 2 m or more set some seeds. The significance of this finding in *S. erectum* which forms large clones is unclear. Although no precise measurements were made it seems that the stigmas remain receptive for at least two days and that wind pollination over distances of several metres or more may occur.

Dispersal

Both Guppy (1906) and Praeger (1913) observed that the fruits of *Sparganium* may remain floating for more than a year under experimental conditions. The fruits are

coated with a hydrophobic material soluble in detergent. Our observations in nature are that after being frozen in ice the exocarp usually ruptures and the spongy mesocarp soon decays in spring liberating the endocarp which sinks. In nature the fruits rarely remain floating for more than 6 months.

Floating fruits may be dispersed by water movements induced by gravity or by wind. Being hydrophobic the fruits readily stick to objects in the water and some fruits are probably dispersed by boats and epizoically on aquatic animals, particularly larger birds. The fruits of *Sparganium* are important food for water birds (Martin & Uhler 1939). In the summary of his work Guppy (1906) stated that seeds that have passed through the alimentary canal of a duck have a higher germination rate; Cook (1962) found no difference in germination rate in seed fed to chickens.

Ecology

Sparganium is found in a wide variety of permanent or seasonally aquatic habitats. The seedlings of all species develop only when submerged in water and thus standing water is necessary for establishment by seed; the rhizomes and corms of some species can, however, perennate terrestrially so that *Sparganium* may inhabit bogs or moors without standing water. In all species the leaves die down in winter and the plants hibernate embedded in the substrate as corms or rhizomes.

The germination and seedling development of *S. erectum* are described and illustrated by Cook (1962). The seed, or more correctly endocarp, is very uniform in structure throughout the genus and is well described by Dietz (1887) and Saccardo (1895). The seed is enclosed in a hard scleridial endocarp with a plugged micropyle. The plug has a somewhat complicated development described by Hegelmaier (1874). Germination only takes place under water or in a fully saturated atmosphere and appears to be fully dependent on dislodging the micropyle or breaking the endocarp. No dormancy has been reported. The first stage in germination consists of an elongation of the cotyledon stalk. This elongation is due to the seed taking up water and can be demonstrated on dead seeds. The first visible indication that the second or vital stage has started is that the tip of the cotyledon stalk turns green and starts swelling as the embryo grows. If seeds without micropylar plugs are left in the dark the cotyledon stalk elongates in a somewhat disorganized spiral without any growth of the embryo. If a seedling is taken from the dark and put into the light there is growth of the embryo and orientation toward gravity. This acquired geotropic response is irrespective of the direction of light and is retained if seedlings are replaced in the dark.

The seedlings of all species of *Sparganium* are very similar and it is not possible to distinguish the species on the basis of the juvenile leaves. *S. natans* develops flowers in the juvenile phase and is thus essentially a submerged species. Other species, such as *S. angustifolium* or *S. gramineum*, develop floating leaves after the submerged juvenile stage; the floating leaves have clearly differentiated ab- and adaxial surfaces. These species flower at the floating-leaved phase. Others, such as *S. erectum* and *S. eurycarpum*, develop erect emergent leaves directly after the submerged juvenile phase. The erect, emergent leaves also show ab- and adaxial differentiation and may be phylogenetically tied to floating leaves. These species flower only at the erect emergent-leaved phase and they are essentially terrestrial plants following their brief aquatic juvenile phase. Most other species occupy positions intermediate between these three extremes (submerged, floating and erect) and show varying degrees of "aquaticness".

Species such as *S. emersum*, *S. fallax* and *S. subglobosum* show considerable plasticity and can flower in floating and emergent phases.

The flexibility in leaf form of some species has enabled them to occupy places subject to rapid changes in water level. The relatively deep roots, rhizomes and stolons also make the plants difficult to dislodge and they are capable of withstanding severe floods and spates. These very phenotypically plastic species are usually poor competitors and rarely build large stands. The less flexible species, either erect-emergent, floating or submerged, are relatively strong competitors and may become dominant over large areas.

The evolutionary trend from small aquatic species to large terrestrial species is clear and is reflected in the order of species in this work. However it is difficult to distinguish between the processes of paedogenesis (sexual maturity in the juvenile phase) and neoteny (retention of juvenile characters in the adult). Although it is popular today to interpret the evolution of flowering plants in terms of reduction, there is no evidence that *S. natans* is more than a neotenous or paedogenetic derivative of *S. eurycarpum*, in terms of life-form it may well be the most primitive.

Parasites and animal feeders

Despite the fact that no comprehensive survey has been attempted, an extraordinary large number of parasites and animal feeders have been recorded on *Sparganium* (13 beetles, 7 moth larvae and 24 fungi). Cook (1962) presented a list based on work done in Britain on *S. erectum* and Müller-Doblies, U. & Müller-Doblies, D. (1977) in Hegi's Flora of Central Europe extended and added to this list.

a) Animal feeders or parasites

Coleoptera. Chrysomelidae: *Donacia appendiculata* Ahr., *D. aquatica* L., *D. bicolor* Zschach, *D. cineria* Hbst., *D. marginata* Hoppe, *D. simplex* F., *D. sparganii* Ahr., *D. tomentosa* Ahr., and *D. vulgaris* Zschach; Cryptophagidae: *Cryptophagus sparganii* St.; Curculionidae: *Thryogenes festucae* Hbst.; Mycetophagidae: *Telmatophilus sparganii* Ahr. (a monotypic genus) and *Typhaea caricis* Ol.

Lepidoptera. Noctuidae: *Archana algae* Esp., *A. sparganii* Esp., *Eustrotia candidula* Schiff., *Phytometra festucae* L.; Pyralididae: *Haemylis sparganiella* Thb., *Hydrocampus stagnata* Don., and *Laelia coenosa* Hb. Several aquatic molluscs, some rodents, ruminants and birds eat the shoots, rhizomes and fruits but none apparently are dependent on *Sparganium*.

b) Plant parasites (including some essentially saprophytic species that are sometimes weakly parasitic)

Chytridiomycetes: *Cladochytrium sparganii-ramosii* Büsgen, and *Urophlyctis ramosa* Büsgen.

Ascomycetes: *Acanthophiobolus helmithosporus* Berl., *Leptosphaeria clava* (Cooke & Auersw.) Sacc., *L. duplex* (Sow.) Sacc., *L. riparia* Sacc., *L. sparsa* (Fuck.) Sacc. var. *meiospora* Feltg., *Metaspheeria sparaganii* Fautrey, *Phaeosphaeria eustoma* (Fuck.) L. Holm, *P. typharum* (Desm.) Karsten, *Pleospora sparganii* Cooke, *Sordaria sparganicola* Phill. & Plowr., *Sphaerella taediosa* Pass., and *S. thais* Sacc.

Deuteromycetes: *Hendersonia sparganii* Niessl., *Macrosporium sparganii* Lindau, *Ramularia sparganii* Lindroth, *Septoria sparganii* Pass., and *Stagonospora sparganii* (Fuck.) Allescher.

Basidiomycetes: Agaricales: *Psathyrella typhae* (Kalchbr.) Pears & Dennis; Tremellales: *Dacryopsis typhae* Höhn; Uredinales: *Uromyces sparganii* Clint & Pech.; Ustilaginales: *Entyloma sparganii* Lagerh., and *Melanotaenium sparganii* Lagerh.

Applied aspects

Sparganium is not of great economic importance. In northwestern Europe *S. erectum* and *S. emersum* are sometimes considered to be undesirable weeds in small rivers, canals or drainage ditches. However, the negative aspects (reduction of water flow and eventual blockage) should be weighed against the benefits, as *Sparganium* may protect the banks from erosion. Also it is a valuable food for wild fowl and offers shelter to numerous aquatic or wetland animals. The Klamath Indians of Oregon used the starchy corms for food but only for subsistence. It is not given important medicinal properties in western Herbals and Pharmacopoeias. In China, according to Duke & Ayensu (1985), it is used to dissolve clots or correct enterrhagia, also as an abortifacient, emmenagogue, lactagogue, sedative and to stimulate blood circulation and to cure cancer of the cervix and liver, lymphosarcoma and relieve abdominal and chest pains. We have no first-hand experience of its efficacy. Although often found in decorative pools it is rarely offered for sale in plant catalogues.

Key to Sparganium

- 1A. Perianth segments more or less translucent, without a thickened dark-brown to black pad of tissue near the apex; margin at apex distinctly erose (see Fig. 2a) subgenus *Xanthosparganium*
- 2A. Distal part of leaves and lower bracts convex or flat but never keeled or triangular in transverse section, usually floating or submerged.
 - 3A. Male heads 1–4, crowded and overlapping at anthesis (appearing as 1 elongated head terminating the main axis); beak of fruit absent or more or less erect, not curved or deflexed, inflorescences simple.
 - 4A. Styles less than 0.5 mm long; stigmas not more than 0.5 mm long; beaks absent or nipple-like 1. *S. hyperboreum*
 - 4B. Styles more than 0.5 mm long; stigmas more than 0.5 mm long; beaks elongate (not nipple-like), at least 0.5 mm long.
 - 5A. Lowermost bract shorter than or scarcely exceeding the inflorescence; all female heads usually axillary 2. *S. natans*
 - 5B. Lowermost bract very much longer than the inflorescence; some female heads usually supraaxillary.
 - 6A. Male heads clearly separated from the uppermost female head; female heads remote (not congested); stigmas usually more than 0.8 mm long; anthers usually more than 0.9 mm long 3. *S. angustifolium*
 - 6B. Male heads contiguous with the uppermost female head; the upper female heads crowded; stigmas not more than 0.8 mm long; anthers usually less than 0.9 mm long. 4. *S. glomeratum*

- 3B. Male heads 2–6, the lower ones remote, not overlapping at anthesis; beak of fruit usually curved or deflexed, inflorescences branched or simple.
- 7A. Stigmas not exceeding 0.8 mm long; beaks not more than 1.5 mm long; fruit shiny, not more than 3 mm long; endocarp not more than 2.5 mm long; leaves usually less than 5 mm wide; inflorescence simple or with one branch (Eurasia) 5. *S. gramineum*
- 7B. Stigmas 1 mm or more long; beaks 2 mm or more long; fruits dull, 3 mm or more long; endocarp 3 mm or more long; leaves usually more than 4 mm wide; inflorescence usually with more than one branch (N. America) 6. *S. fluctuans*
- 2B. Distal part of the leaves and lower bracts keeled or triangular in transverse section, usually erect and emergent.
- 8A. Male heads 1–2, when 2 then overlapping at anthesis; lowermost male head contiguous with uppermost female head (without a sterile internode); stigmas up to 0.8 mm long 4. *S. glomeratum*
- 8B. Male heads 3–10, at least the lower one remote at anthesis; lowermost male head separated from uppermost female head by a sterile internode; stigmas more than 0.8 mm long.
- 9A. Female heads usually remote, the lower often pedunculate; lower bracts shorter or about equalling the inflorescence; fruit beak shorter than the fruit body in length; fruit body 3.5–5.5 mm long 7a. *S. emersum* subsp. *emersum*
- 9B. Female heads crowded and usually sessile (the lowermost sometimes remote and pedunculate); lower bracts conspicuously longer than the inflorescence; fruit beak equal to or exceeding the fruit body in length; fruit body 3–4 mm long (N.E. America) 7b. *S. emersum* subsp. *acaule*
- 1B. Perianth segments not translucent, with a thickened darkbrown to black pad of tissue near the apex; margin at apex emarginate to entire or nearly so (see Fig. 2a) subgenus *Sparganium*
- 10A. Stigmas less than 1.5 mm long; inflorescence simple or branched, when branched rarely more than 2 branches bear female heads; endocarps more or less smooth *S. americanum* group
- 11A. At least 2 female heads supra-axillary and at least one concaulescent up to or beyond the next internode (inflorescence simple; female heads widely spaced) 8. *S. fallax*
- 11B. Female heads axillary or on lateral branches, occasionally 1 head supra-axillary but never concaulescent to next internode (inflorescence simple or branched; female heads spaced or upper approximate).
- 12A. Fruit obovoid to almost globose, shiny and light brown and below, subsessile or with an up to 1 mm long pedicel; endocarp rarely more than 3 mm long; stigma rarely more than 1 mm long; lowermost bract shorter or as long as the inflorescence (inflorescence usually branched, lowest branches bearing both male and female heads or male heads only) (E. Asia, Australasia) 9. *S. subglobosum*
- 12B. Fruit ellipsoidal or fusiform, shiny or dull above and dull below, with pedicels more than 1 mm long; endocarp 3.5 mm or more long; stigma usually more than 1 mm long; lowermost bract very much longer than the inflorescence (inflorescence branched or simple).

- 13A. Fruits less than 5 mm long with beaks less than 4.5 mm long, dull and dark-brown above below; fruiting heads less than 25 mm diameter; endocarp 4 mm or less long.
- 14A. Upper female heads approximate; fruit without dark brown glands, not pitted; beak not more than 4 mm long; pedicels usually 3 mm or more long; inflorescence never branched (E. Asia) 10. *S. japonicum*
- 14B. Upper female heads remote; fruit with dark-brown glands and pits; beak 3 mm or more long; inflorescence simple or branched (when branched the branch or branches bearing both male and female heads), (N. America) 11. *S. americanum*
- 13B. Fruits more than 5 mm long with beaks more than 4 mm long, shiny and light-brown above, dull and pitted below; fruiting heads 25 mm or more in diameter; endocarp more than 4.5 mm long (inflorescence simple or with lowermost branch bearing male heads only; anthers 1.0–1.6 mm long) (N. America) 12. *S. androcladum*
- 10B. Stigmas more than 1.5 mm long; inflorescence usually branched with more than 2 branches bearing female heads, if less then endocarps with longitudinal ribs.
- 15A. At least some female heads borne sessile on main axis; endocarps more or less smooth with shallow longitudinal furrows but not ribbed; fruits with 1–4 mm long pedicels; female heads axillary or supra-axillary.
- 16A. Fruiting heads less than 25 mm diameter: fruits rarely more than 5 mm long, dull and pitted above and below; stigmas rarely more than 2 mm long; lowermost branch bearing both male and female heads 11. *S. americanum*
- 16B. Fruiting heads more than 25 mm diameter; fruits more than 5 mm long, shiny above and dull, pitted and glandular below; stigmas more than 2 mm long; lowermost branch bearing male heads only 12. *S. androcladum*
- 15B. Female heads borne on axillary branches or peduncles, very rarely sessile on main axis but if so then axillary; endocarps with longitudinal ribs; fruits sessile or pedicels up to 1.5 mm long *S. erectum* group
- 17A. Less than half the ovaries bilocular; less than half the stigmas bifid *S. erectum* sensu lato
- 18A. Fruits obpyramidal, flattened above, contracted very abruptly into a beak; viewed from above distinctly angled 13a. *S. erectum* subsp. *erectum*
- 18B. Fruits fusiform to spherical, domed to conical above, tapering into a beak, viewed from above not or indistinctly angled.
- 19A. Fruits with distinct shoulders, upper and lower parts differing in form, colour and texture.
- 20A. Upper part of the fruit light-brown to straw-coloured, conical to domed but not distinctly wider than lower part; slightly constricted below the shoulder; fruits sessile or subsessile 13b. *S. erectum* subsp. *stoloniferum*

- 20B. Upper part of fruit brown to black, matt or shiny near the shoulder, domed and inflated, wider than the lower part, distinctly constricted below the shoulder; fruits with pedicels up to 1.5 mm long 13c. *S. erectum* subsp. *microcarpum*
- 19B. Fruits without distinct shoulders, upper and lower parts alike in form, colour and texture.
- 21A. Fruits ellipsoidal to fusiform, gradually tapering into a beak; beaks at least 2 mm long 13d. *S. erectum* subsp. *neglectum*
- 21B. Fruits widely ovoid to almost spherical, abruptly tapering into a beak; beaks less than 2 mm long 13e. *S. erectum* subsp. *oocarpum*
- 17B. More than half the ovaries bilocular; more than half the stigmas bifid 14. *S. eurycarpum* sensu lato
- 22A. Fruits flattened above; fruiting heads 15–32 mm diameter; endocarps 7–10 mm long (N. America) 14a. *S. eurycarpum* subsp. *eurycarpum*
- 22B. Fruits domed or pyramidal above; fruiting heads rarely exceeding 20 mm diameter; endocarps 6–7 mm long (N.E. Asia) 14b. *S. eurycarpum* subsp. *coreanum*

Species of subgenus Xanthosparganium

1. ***Sparganium hyperboreum*** Beurling ex Laestadius, Wikström's Årsberättelse, 1850, Bihang: 4. 1853 or 1854; in Beurling, Öfversigt Kongl. Vetenskaps-Akad. Förhandl. (Stockholm), 9 (8): 192. 13 October 1852 the names are cited without description using "Laest. in schaed". Type: Sweden, Torne Lappmark, *Laestadius* (lectotype: chosen here, S, sheet bearing "*S. hyperboreum* Laest. var. *natans*, Kengis, Laestadius"; a possible isolectotype in S is dated 1842 by Laestadius; see Cook, Bot. Jahrb. Syst., 107: 275. 1985 for details of typification).

= *S. natans* var. *submuticum* C. J. Hartman, Handbok Skand. Flora, ed. 4: 312. October 1848 ≡ *S. submuticum* (C. J. Hartman) L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Flora, ed. 12: 108. October 1889. Type: lectotype to be chosen from material in LD.

= *S. submuticum* forma *platyphylla* [sic] L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Flora, ed. 12, 108. October 1889. Type: lectotype to be chosen from material in LD.

= *S. williamsii* Rydberg, N. Amer. Flora, 17: 10. 30 June 1909. Type: Canada, Yukon Terr., Klondyke Bottom, 9 & 23 July 1899. R. S. Williams (holotype: NY).

Slender plants with vegetative parts submerged, or floating, or in summer sometimes emergent. Stolons short, up to ca. 10 cm long and ca. 0.8–1.2 mm diameter. Scale leaves up to 9 mm long and 1 mm wide.

Basal leaves usually floating or emergent, relatively thick, opaque, yellowish-green, (5–)10–40(–80) cm long and (0.5–)1–3(–5) mm wide, flat, without a distinct midrib and without a keel even at the base, gas chambers in one layer sometimes higher than wide; sheaths not inflated; apex rounded, not thickened.




Fig. 5. Diagrammatic representation of inflorescence of: a, *Sparganium hyperboreum*; b, *S. hyperboreum* \times *S. natans*; c, *S. natans*.

Flowering stems (5–)10–30(–60) cm long, 0.8–2 mm diameter, with rarely more than 4 elongated sterile nodes below the inflorescence, decumbent or ascending, usually floating, flexuous or rarely erect and emergent.

Inflorescence bracts with very slightly inflated bases and hyaline margins; the lowermost bract 2–6(–14) cm long, distinctly exceeding the inflorescence (Fig. 5a). Uppermost bract absent in fruiting plants.

Inflorescence simple, 2.5–4.5 cm long.

Female heads (1–)2–3(–4), usually congested or adjacent (not remote), the lower one or two usually supra-axillary with peduncles adnate below to the main axis (see Fig. 5a); the peduncle of the lowest head mostly exceeding 5 mm, often arched; in fruit (5–)7–11(–14) mm diameter.

Male head terminal, solitary, (or 1 or 2 contiguous heads appearing as 1), adjacent to the uppermost female head or with an internode less than 5 mm long.

Male flowers with filaments 4–5 mm long; anthers oblong, (0.3–)0.4–0.8 mm long.

Female flowers with perianth segments slender, linear-spathulate, with erose tips, one third to one half but rarely more as long as the fruit, attached at the base of the fruit or along a short pedicel; stigmas 0.3–0.5 mm long, elliptic to almost orbicular, obliquely attached to a less than 0.5 mm long style.

Fruits ellipsoid to ovoid-fusiform, (1.5–)3–4(–5) mm long, (1–)1.5–2.5 mm diameter, often constricted around the middle, brown or yellowish, dull-surfaced, tapering below to an obconic base with short pedicel, rather abruptly tapered distally to an acute or rounded summit with a minute beak, 0–0.5 mm long (see Fig. 6a); endocarp widely ellipsoidal often rounded with a somewhat asymmetrical apex and elongate but




Fig. 6. Perianth segments, fruits and endocarps of: a, *Sparganium hyperboreum*; b, *S. natans*; c, *S. angustifolium*; d, *S. glomeratum* (scale bar: 1 mm).

blunt base; 2–2.5 mm long, 1.–1.7 mm diameter, smooth or with few, very fine, longitudinal furrows.

Diagnostic features

It can be distinguished from all other species of *Sparganium* by its fruit which has a minute, nipple-like beak. It is patristically related to *S. natans* with which it occasionally hybridizes (for description of hybrid see p. 237 and Table 2).

Distribution

It is found in arctic and boreal N. America and Eurasia. It extends from Scandinavia eastwards through Russia and Siberia to Kamchatka and Hokkaido. In N. America it reaches from the Pacific coastal forests of the Yukon into Alaska and across to the Atlantic coast extending southwards to S. Newfoundland, Anticosti Island and Cape Breton Island. It is also found in S. W. Greenland and Iceland. It is almost confined to the Taiga but occasionally extends into the Tundra and is known from a small region in the southern Alps in N. Italy, (see Maps 1 & 2). In the Alps it grows at an alti-

tude of between 2250 m and 2350 m elsewhere it is rarely found above 1000 m. Fossil fruits have been reported from Poland so it is likely that the alpine populations represent genuine glacial relicts.

Ecology

Unfortunately, we have not had the opportunity to study living material of *S. hyperboreum*. According to the literature it is found in flat-bottomed pools and ditches in the Taiga, Tundra and high Alpine Zones, preferring cold water. Glück (1938) gives description of the ecology of the alpine populations. It appears to be rather indifferent to water quality being reported in calcium rich and calcium poor water; however, it is found in neither very oligotrophic nor highly eutrophic conditions.

It is essentially a plant of shallow water but grows at depths of up to about 80 cm but is rarely found in flowing water and is not resistant to much wave action. The leaves are usually completely submerged or partly floating but unlike *S. natans* it apparently rarely develops truly aerial leaves and rarely flowers in a terrestrial state. The leaves die in autumn and it overwinters as a corm. It is found growing with species such as *Utricularia ochroleuca* Hartman, *Subularia aquatica* L., *Hippuris vulgaris* L., *Potamogeton alpinus* Balbis, *P. filiformis* Persoon, *P. gramineus* L. and *Ranunculus trichophyllum* subsp. *eradicatus* (Laest.) C. D. K. Cook.

Variation

The variation is well described by Harms (1973) and the pattern he describes for N. America is similar in the Old World except that hybridization is local and hybrids are rare. We agree that *S. hyperboreum* occasionally hybridizes with *S. natans* (See p. 237). We are not, however, convinced that introgressive hybridization takes place although perhaps occasionally back crosses are found. *S. hyperboreum* has been recorded to make hybrids with *S. angustifolium* (e.g. Lid 1952, Benum 1958) and *S. emersum* (Lid 1952). There is no experimental evidence and we do not find the morphological evidence convincing.

2. ***Sparganium natans* L.**, Species Plantarum, 971. 1753. Type: "in Europae borealis lacubus, paludibus" (lectotype: Linn. Savage Cat. No. 1095.2; see Cook, Bot. Jahrb. Syst., 107: 272. 1985); non L., Flora Lapponica, 345.1737, nec L., Flora Suecica, 771. 1745.
= *S. minimum* J. Ray, Historia Plantarum, 1910. 1688; nom. illeg., pre-1753 (cited by Linneus as a synonym of *S. natans*) ≡ *S. minimum* J. Hill, British Herbal, 507. 1756; nom. illeg., binomial names not used consistently (ICBN Art. 23) ≡ *S. natans* L. var. *minimum* C. J. Hartman, Handbok Skand. Flora, ed. 1: 43. Oct. 1820 ≡ *S. minimum* Wallroth, Erster Beitrag Fl. Hercyn., 2: 294. 1840 ≡ *S. minimum* E. M. Fries, Summa Veg. Skand., 2: 560. 1849; based on J. Ray. Type: the description in J. Ray, Historia Plantarum, 1910. 1688.
= *S. natans* var. *subdecumbens* Laestadius, Wikström's Årsberättelse, 1850, Bihang: 4. 1853 or 1854. Type: Sweden, "lacu Rättar-Dammen prope Drottningholm"? *Laestadius* (holo- or lectotype: S, n.v.)
= *S. natans* var. *suberectum* Beurling ex Laestadius, Wikström's Årsberättelse, 1850, Bihang: 4. 1853 or 1854. Type: Sweden, "in Lapponica Umensi, Dr. Ångström", (holo- or lectotype: S?)

- = *S. rostratum* Larsson, Fl. Wermland, 260. 1859 ≡ *S. minimum* E. M. Fries forma *rostrata* [sic] (Larsson) L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Flora, ed. 12, 108. October 1889 ≡ *S. minimum* E. M. Fries var. *flaccidum* subvar. *rostratum* (Larsson) Graebner in Engler, Pflanzenreich, 2 (IV. 10): 23. 1900. Type: Sweden, specimen not yet located.
- = *S. ratis* Meinshausen, Bull. Soc. Imp. Nat. Moscou, N.S., 3: 174. 1890; emended in Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4: 39. December 1893 ≡ *S. minimum* E. M. Fries var. *oligocarpon* (Ångström) subvar. *ratis* (Meinshausen) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora, 1: 293. 15 June 1897; Type: USSR, "Ingrien kleiner Waldseen im Nordgebiete, July" coll.? Meinshausen (lectotype: chosen here, the specimen labelled "Typus" by N. Tzvelev in 1958: LE; isolectotypes: L, MW).
- = *S. septentrionale* Meinshausen, Bull. Soc. Imp. Nat. Moscou, N.S., 3: 174. 1890; emended in Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4: 39. December 1893 ≡ *S. minimum* E. M. Fries var. *oligocarpon* (Ångström) subvar. *septentrionale* (Meinshausen) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora, 1: 293. 15 June 1897. Type: USSR, "Ingrien, nur im Nordgebiete in rieselten kalten Quellbächen der Torfmoore" "mit *Malaxis paludosa* nicht häufig" (lectotype: chosen here, specimen with "in paludibus Malaxide paludosae, rarius. Julio" coll.? Meinshausen, the specimen labelled "Typus" by N. Tzvelev in 1958: LE; isolectotypes: LD, MW).
- = *S. flaccidum* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4: 37. December 1893 ≡ *S. minimum* E. M. Fries var. *flaccidum* (Meinshausen) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 292. 15 June 1897. Type: USSR, "Ingrien in tieferen Sumpfen mit faulenden, dunklen Gewässern untergetaucht" (lectotype: chosen here, specimen with "in fossis profundis aquis putredis submersum, Jul." Meinshausen with "N12/2296A" written on the sheet: LE; isolectotypes: LD, MW).
- = *S. perpusillum* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg, N.S. 4: 38. December 1893 ≡ *S. minimum* E. M. Fries var. *perpusillum* (Meinshausen) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 293. 15 June 1897. Type: USSR, "Ingrien" "am Gestade des finnischen Meerbusens, im Nordgebiete (ipse)" (lectotype: chosen here, the specimen labelled "Lachtam" collected 9 July 1891 by Meinshausen: LE; isolectotype: LD).
- *S. minimum* var. *strictum* (distinct from var. *typicum*) Luersson ex Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 293. 15 June 1897, based on Luersson, Schrift. Phys.-ökön. Ges. Königsberg, 29: 59. 1888 und 1889, nomen nudum. Type: Poland, "Ostpreußen Gutten bei Johannisburg, Luersson (holotype: destroyed in B?).

Slender plants with vegetative parts usually submerged or floating. Stolons up to ca. 20 cm long and 1–2 mm diameter. Scale leaves up to 12 mm long and 2 mm wide.

Basal leaves when submerged or floating, thin, translucent, deep green, 6–40(–60) cm long and (1.5–)2–6(–10) mm wide (or when terrestrial 7–20 cm long and 2.5–4 mm wide), flat to somewhat concave but without a distinct midrib and without a keel even at the base, gas chambers in one layer and wider than high; sheaths not inflated; apex rounded, not thickened.

Flowering stems (6–)8–40(–100) cm long, 1–3 mm diameter, with (4–)5–8(–9) elongated sterile nodes below the inflorescence, decumbent or ascending, usually floating, flexuous or rarely erect and emergent.

Inflorescence bracts with usually somewhat inflated bases with hyaline margins; the lowermost bract 1–5(–8) cm long, scarcely exceeding the inflorescence (Fig. 5c), the uppermost leaf-like or scale-like.

Inflorescence simple, 1.5–8 cm long.

Female heads 1–3(–4), usually remote, axillary or rarely supra-axillary; the uppermost sessile, the lowermost sessile or with an up to 28 mm long but usually less, more or less straight peduncle; in fruit (5–)7–10(–15) mm diameter.

Male head terminal, solitary, (of 1 or 2 contiguous heads appearing as 1, remote from the uppermost female head, internode exceeding 5 mm long.

Male flowers with filaments 3–5 mm long; anthers oblong, (0.3–)0.4–0.5(–0.8) mm long.

Female flowers with perianth segments elliptic to cuneate-spathulate with erose tips, scarcely clawed below, one half to two thirds as long as the fruit, attached at the base of the fruit or along a short pedicel; stigmas (0.3–)0.5–0.8 mm long, ovate, obliquely attached to a 0.5–1.5 mm long style.

Fruits ellipsoid to narrowly obovoid-fusiform, (1.5–)2–4(–6) mm long, 1.0–2.5 mm diameter, only slightly if at all contracted near the centre, greenish or brownish, dull-surfaced, tapering below, sessile or with a pedicel not exceeding 1 mm long, rather abruptly tapered distally to an acute summit with a conic based, slender beak, 0.5–1.0(–1.5) mm long; endocarp ovoid, 2–2.5 mm long, 1.2–1.4 mm wide, uniformly narrowed below, smooth or with very fine longitudinal furrows (Fig. 6b).

Diagnostic features

Inflorescence simple; female heads axillary; male head solitary and remote; leaves flat, thin, translucent and dark-green; the lowest bract scarcely longer than the inflorescence. Like *S. hyperboreum* but female heads remote and separated from the male head by an at least 5 mm long internode; fruit with a slender beak. See Table 2 for further information.

Distribution

It is found in arctic and boreal N. America and Eurasia. It extends from Newfoundland to Alaska, south (on higher ground) into Oregon, Utah, Colorado and New Jersey. It is characteristic of Great Lakes conifer-hardwood and western subalpine-montane forest regions. Across much of N. America the northern limit of *S. natans* coincides well with the northern limit of trees (Harms 1973).

Table 2. Comparison of *Sparganium hyperboreum*, *S. Natans* and their hybrid.

Character	<i>S. hyperboreum</i>	Hybrid	<i>S. natans</i>
Leaf width	1–3 (–5) mm	2–5 mm	2–6 (–10) mm
Leaf thickness	Thick, opaque	Thick, opaque	Thin, translucent
Leaf colour	Yellowish	Light-green to brown	Dark-green
Female heads dispersion	Congested above	Remote	Remote
Female peduncle adnation	Adnate	Adnate	Axillary
Female lowest peduncle length	> 5 mm	3–7 mm	< 3–8 mm
Distance between ♂ and ♀ heads	< 5 mm	4–10 mm	> 5 mm
Fruit colour	Yellowish	Light-green to brown	Dark-green
Fruit beak length	< 0.5 mm	0.5–1.0 mm	About 1 mm

In Europe it extends from southern Scandinavia (with isolated stations above 65° N) southwards to the Pyrenees, N. Appennini and S. Bulgaria. In Scandinavia it is found up to almost 1000 m and in the Alps up to about 2300 m. In Asia it extends across Siberia from Yenisei (66° N) south to Omsk. It is also recorded from the Caucasus, Sajan Mountains and northern Japan, see Maps 3 & 4. The Japanese record is based on a single specimen: 'Azuma', 13 July 1893. *Faurie* 10368 (K).

Ecology

S. natans is found in sheltered bays or inlets of lakes, in pools, ditches and drainage channels. It is a characteristic species of peat diggings, is rarely found in flowing water and is not resistant to much wave action.

It is essentially a species of shallow water rarely being found more than 60 m deep, although it has been recorded at depths of more than 1 m. It flowers best in water 20–40 cm deep. In deeper water it develops fewer inflorescences and each inflorescence tends to have fewer heads. Both leaf surfaces are wettable and it does not develop true floating leaves. The upper leaves and inflorescence bracts are sometimes aerial. In summer *S. natans* can develop terrestrially but the terrestrial state is a rather shy flowerer and it is readily killed by excessive drought or frost and therefore rarely develops ripe fruit. The aerially developed leaves are shorter, more glossy and deeper green than the submerged ones.

The species shows a preference for slightly acid, mesotrophic to somewhat oligotrophic water with a base-rich substrate. It has been observed to tolerate salinities of up to 0.22% in Finland. It is usually found in peat-rich muds preferring aerobic conditions (Gyttja) and rarely grows in clay or sandy soils.

It usually grows in rather loose stands and is rarely dominant over large areas although it is the characteristic species of the association "Sparganietum minimi" within the Class "Utricularietae intermedio-minoris". The leaves die in winter and it overwinters as a small corm. The stolons rarely develop more than 3 to 5 internodes in a year; it is likely that the effective spread is through seeds.

In open water *S. natans* is frequently found growing with *Potamogeton* species, such as *P. polygonifolius* Pourret & Figeac, *P. alpinus* Balbis, *P. natans* L. and *P. gramineus* L. It often grows in loose stands of *Phragmites* and *Carex rostrata* Stokes between the upright stems. It sometimes grows with *S. angustifolium* but rarely grows intermingled as it occupies the shallower water or shaded parts of pools. In similar pools without *S. natans*, *S. angustifolium* is capable of growing in shallow water and shade; when *S. angustifolium* is absent *S. natans* can invade the deeper water. It is also occasionally found growing with *S. emersum*.

Notes

S. natans has been reported to hybridize with *S. angustifolium* (Lid 1952), *S. emersum* (see p. 255) and *S. hyperboreum* (Lid 1952). None of these hybrids have been experimentally verified. From morphological and ecological evidence we are convinced that *S. natans* × *S. hyperboreum* exists; the two other putative hybrids are doubtful and we are not convinced that they exist.

Hybrid

1 × 2. *Sparganium hyperboreum* × *S. natans*. This hybrid is like *S. hyperboreum* in having supra-axillary female heads but distinguished by wider (2–5 mm) leaves and

green-brown fruits with beaks ca. 0.5–1.0 mm long. The upper (distal) female heads are often more remote than in “pure” *S. hyperboreum* and the internode between the uppermost female and the male head can be up to ca. 10 mm long (Fig. 5 b). The combination of supra-axillary female heads and beaked fruits is characteristic (see Table 2).

Despite hybridization, each of the parents retain their essential integrity and pose few taxonomic problems. The hybrid is infrequent through much of the world (see Maps 3 & 4); the distribution of the parents is mostly allopatric. The main exception is in C. Alaska and N.W. Canada where this distinction breaks down (probably due to the topography) and putative hybrids are found. In the Old World hybrids are much rarer. They have been recorded from Bygland and Anot in Norway by Lid (1959) and from Troms Fylke by Benum (1958).

The hybrid appears to be fertile. Harms (1973) suggests that introgression may occur but we feel that he is using the term rather loosely. Some backcrossing or segregation may occur but most hybrids look like first generation progeny.

3. ***Sparganium angustifolium*** Michaux, Flora Boreali-Americanæ, 2: 189. 1803 ≡ *S. natans* var. *angustifolium* (Michaux) Pursh, Flora Amer. Septentrionale, 1: 34. December 1813 ≡ *S. simplex* var. *angustifolium* (Michaux) Engelmann in Gray, A., Manual Bot. Northern U.S., ed. 5, 481. 1867 ≡ *S. emersum* var. *angustifolium* (Michaux) R. L. Taylor & B. McBryde. Canad. J. Bot., 56: 193. 1978. Type: Canada, Lake Mistassini, Michaux (holotype: P n.v.).
- = *Isoetes lacustris* var. *fluitans* Döll, Rheinische Flora, 40. June 1843. Type: none found, perhaps in KR.
- = *Sparganium affine* A. Schnizlein, Die Pflanzenfamilie der Typhaceen (Nordlingen), 27. 1845. Type: lectotype to be found perhaps in ER or REG.
- = *S. oligocarpon* Ångström, Bot. Not. 1853: 149. 1853. ≡ *S. minimum* E. M. Fries var. *oligocarpon* (Ångström) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora, 1: 292. 1897. Type: Sweden, “at Lycksele et Betsele Lapponiae Umensis”, July., Ångström (lectotype to be chosen in S, isolectotypes: S. UPS and in other herbaria).
- = *S. vaginatum* L. M. Larsson, Flora Wermland och Dal, 259. 1859. Type: Sweden, “Wermland, vester om Liljendal i Bergsladen” L. M. Larsson (holotype: C, n.v.).
- = *S. borderi* [sic] W. O. Focke, Abhandl. Naturwiss. Ver. Bremen, 5: 409. April 1877; ≡ *S. affine* subsp. *borderi* (W. O. Focke) Weberbauer ex Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, I: 288. 15 June 1897; (Graebner clearly intended it to be a subspecies, the “B” is misprinted in roman and should have been put in italic). Type: France, “Hautes Pyrénées, Tremouse”, *Bordère* (holotype: BREM n.v.; topotypes: “Tremouse, July 1886, *Bordère*” B, Z).
- = *S. simplex* var. *multipedunculata* [sic] Morong, Bull. Torrey Bot. Club, 15: 79. 2 March 1888 ≡ *S. simplex* proles *longissimum* var. *multipedunculatum* (Morong) Graebner in Engler, Pflanzenreich, 2 (IV, 10): 17. September 1900 ≡ *S. multipedunculatum* (Morong) Rydberg, Bull. Torrey Bot. Club, 32: 598, 1905 ≡ *S. emersum* var. *multipedunculatum* (Morong) Reveal, Taxon, 19: 797. 1970 ≡ *S. angustifolium* subsp. *emersum* var. *multipedunculatum* (Morong) Brayshaw, Occasional Papers British Columbia Provincial Museum, 26: 115. 1985. Type: USA, Montana, Great Falls, s.d., R. S. Williams (holotype: NY).
- = *S. affine* var. *zosteraefolium* [sic] L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Flora, ed. 12, 110. October 1889 ≡ *S. affine* subvar. *zosterifolium*

- [sic] (L. M. Hartman) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 288. 15 June 1897. Type: none cited, Sweden, Medelpad Prov., Sundsvall par., 20–25 September 1886. L. M. Neumann (holotype: UPS).
- = *S. affine* var. *deminutum* L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Flora, ed. 12, 110. October 1889 ≡ *S. affine* subsp. *borderi* [sic] var. *deminutum* (L. M. Neuman) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 289. 15 June 1897. Type: "Ligger i Linnés herb. under namnet *Sp. natans* β" it is not known which specimen Neuman saw; *S. natans* var. *β* L. was based on *S. minimum* Ray (1688), Neuman is probably referring to *S. angustifolium* Michaux from his description.
- = *S. affine* var. *microcephalum* L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Flora, ed. 12, 110. October 1889 ≡ *S. affine* subsp. *borderi* [sic] var. *microcephalum* (L. M. Neuman) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 289. 15 June 1897. Type: Plate 1, Fig. 1B & 1C in Bot. Not. 1852.
- = *S. subvaginatum* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4 (36): 34. December 1893, pro parte ≡ *S. simplex* var. *angustifolium* subvar. *subvaginatum* (Meinshausen) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 285. 15 June 1897 ≡ *S. simplex* var. *subvaginatum* (Meinshausen) Graebner in Engler, Pflanzenreich, 2 (IV. 10): 17. September 1900 ≡ *S. emersum* forma δ *subvaginatum* (Meinshausen) Soó, Acta Bot. Acad. Sci. Hung., 17: 124. dated 1971 published 1972. Type: "Finnland, Archipelago Aboensi-Krånskor", pr. Kelo, Ruprecht; Ins. Sitcha, Mertens; Rocky-Mountains, reg. alpina, C. C. Parry (lectotype to be chosen).
- = *S. kawakamii* Hara, J. Jap. Bot., 14: 53, 54. January 1938. Type: USSR, Kurilskiye Ostrova, "Ins. Etorofu, Ponto, Rubetsu, 25 August 1898, Kawakami (holotype: MAK, 13 8613; ? isotype: "Herb. Agr. Hokk. Imp. Univ.")".

Slender plants with vegetative parts usually submerged or floating rarely emergent but when so then very small. Stolons up to ca. 35 cm long and ca. 3 mm diameter. Scale leaves up to 16 mm long and 3 mm wide.

Basal leaves usually submerged below and floating distally, (20–)30–80(–250) cm long, (1.5–)2–4(–10) mm wide, flat or rounded abaxially, without a distinct midrib and without a keel even at the base; abaxial surface of floating portion of leaf with one layer of 6–8 gas chambers, the submerged portion with 2 or rarely 3 layers of gas chambers, the sheath with 2 or 3 layers.

Flowering stems (7–)30–100(–175) cm long, 1–3 mm diameter, with 2–4 elongated sterile nodes below the inflorescence, decumbent or ascending, usually floating, flexuous or rarely erect and emergent.

Inflorescence bracts inflated at base with hyaline margins; lowermost bract (5–)10–20(–60) cm long, (1.5–)2.5–4.0(–4.5) times as long as inflorescence (Fig. 7a); the bract of the uppermost female head 1.5–20 cm long and usually exceeding the inflorescence, the bract of the lowest male head often green and leaf-like. Inflorescence simple, 1–9 cm long.

Female heads (1–)2–4(–5), remote, the lowest axillary or supra-axillary and peduncled the other axillary or supra-axillary and usually sessile; the free part of the peduncle 5–25(–55) mm long; in fruit (8–)10–20(–24) mm diameter.

Male heads (1–)2–3(–4), usually crowded and appearing as one elongated head terminating the main axis (Fig. 7a), usually remote from the uppermost female head with an internode (0–)10–20(–50) mm long.

Female flowers with perianth segments spathulate, translucent, with erose tips, scarcely clawed below, about two thirds as long as the fruit, attached at the base of the




Fig. 7. Diagrammatic representation of inflorescences of: a, *Sparganium angustifolium*; b, *S. glomeratum*; c, *S. gramineum*; d, *S. fluctuans* (scale bar: 1 mm)

fruit or along a short pedicel; stigmas (0.6–)0.8–1.0(–1.2) mm long, lanceolate-ovate, obliquely attached to a 1.5–2 mm long style.

Male flowers with filaments 3–5 mm long; anthers narrowly oblong, (0.8–)0.9–1.0(–1.2) mm long.

Fruits ellipsoid to fusiform, 3–5.5 mm long and 2–2.5 mm diameter, brownish, usually somewhat constricted near the centre, tapering below to an obconic base with a (0.8–)1.0–1.5 mm long pedicel, tapered distally to an acute summit with a conic-based, 1.5–2.0(–2.2) mm long beak; endocarp ovoid, 2.5–4.0 mm long, ca. 2.5 mm diameter, tapered below into a ca. 2.5 mm long stalk, smooth with some fine longitudinal furrows (Fig. 6 c).

Diagnostic features

Inflorescence simple; at least some female heads supra-axillary; male heads up to 3, continuous, clearly separated from uppermost female head, at anthesis appearing as 1 elongated head; stigmas upto 1 mm long; lowest inflorescence bract inflated at base, usually at least twice as long as the inflorescence; leaves usually floating, plano-convex.

Distribution

S. angustifolium is a holarctic species with a circumboreal distribution. It is commoner in suboceanic regions and is rare (or perhaps poorly collected) in continental Eurasia. It shows a somewhat arctic-alpine distribution pattern in Europe and N. America (see Maps 5 & 6).

In N. America it extends from the Aleutian Islands and Alaska to Newfoundland, reaching southwards to California, Idaho, Montana, Wyoming and Utah on higher ground, growing at about 3000 m in Colorado; only in western N. America does it extend beyond 60° N but even then rarely grows beyond the arctic circle. It is present in Greenland and Iceland. In Europe it is widespread and common in the North, extending beyond the arctic circle (reaching 71° N in Norway). In central and southern Europe it is confined to higher land in the Alps and the mountains of Portugal, Spain (2000 m in Sierra de Gredos) and Macedonia. In European Russia it is found in Karelia-Lapland, Dvina-Pechora, Ladoga-Il'nen and the Upper Volga. In eastern Asia it is found in Kamchatka, Sakhalin southwards to Hokkaido and the mountains of Honshu.

Ecology

S. angustifolium is confined to and can be used as an indicator of acid oligotrophic water. It occupies a variety of different habitats from large lakes to small pools, ditches and streams. Among the species of *Sparganium* it is the most tolerant of deep water usually being found in 30–150 cm of water but occasionally growing to 2.5 m or more. It develops true floating leaves with wettable abaxial and unwettable adaxial surfaces. These floating leaves are plano-convex in transverse section with considerable aerenchyma and 6–8 gas chambers. The stems are relatively thin, flexible and non-buoyant. The sheaths of the cauline leaves and inflorescence bracts are often inflated and filled with gas; these buoyant sheaths probably play an important role in keeping the inflorescence above the surface of the water. Usually most of the photosynthetic surface is exposed to air (adaxial surface of floating leaves). In summer *S. angustifolium* is capable of growing terrestrially with short, weakly erect and unwettable leaves.

S. angustifolium can spread by stolons and is sometimes locally dominant with the floating leaves covering quite large areas. Within these dense stands very few other species are found; the commonest associated species outside the dense stands are:

Isoetes lacustris L., *I. echinospora* Durieu or *I. macrospora* Durieu, *Potamogeton alpinus* Balbis, *Littorella uniflora* (L.) Ascherson or *L. americana* Fernald, *Scirpus (Eleogiton) fluitans* L. and *Lobelia dortmanna* L. It is sometimes found growing with *Sparganium natans* and *S. glomeratum*, it apparently sometimes hybridizes with the latter.

In autumn the leaves of *S. angustifolium* die and it overwinters as corms. Local spread is by stolons and perhaps also by seed. It is often infected with the larvae of *Hydrocampus stagnata* (a small moth: Pyralidae) which may well reduce its vigour; it is readily eaten by many other moth and beetle species (see p. 227).

Variation

S. angustifolium is a relatively invariable and constant species. In deep water the lower female heads are usually borne on long peduncles (Fig. 7 a) with highly inflated bract-bases. In very shallow water or on land the female heads are usually sessile and the bract-bases are not inflated.

This variation is most likely to be phenotypically induced but has led to some taxonomic confusion. Deep water plants occasionally carry the epithets "multipedunculatum" or "subvaginatum" while those having the shallow-water or terrestrial state are often called "*S. borderei*". Fertile hybrids are formed with *S. emersum* (see p. 255) and *S. gramineum* (p. 247) which in some regions may lead to taxonomic difficulties.

Hybrids

S. angustifolium has been reported to hybridize with *S. hyperboreum* (Lid 1952; Benum 1958), *S. natans* (Rothert 1910, Lid 1952) and *S. glomeratum* (Rothert 1913, Lid 1952). Although these hybrids are likely we are not convinced they exist and material determined as hybrid is usually attributable to one or other of the putative parents. Hybrids with *S. emersum* (p. 255) and *S. gramineum* (p. 247) do exist and are fertile.

4. ***Sparganium glomeratum*** (Beurling ex Laestadius) L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Flora, ed. 12, 111. October 1889 ≡ *S. erectum* var. *γ glomeratum* Beurling ex Laestadius, Bihang till Wikström's Årberättelse, 1850: 2. 1853 or 1854. Type: Sweden, Hernösand, 1843, Laestadius (lectotype: chosen here, sheet with "*S. glomeratum*" in Beurling's own handwriting and "*S. simplex* var. *glomeratum*" in Laestadius' handwriting: S; isolectotype: S). In Beurling's publication in "Översigt Kongl. Vetenskaps-Acad. Förhand (Stockholm), 9 (8): 192. 13 October 1852" the epithet "*glomeratum*" is a nomen nudum citing "Laest. in schaed" for further details see Cook (1985).
- =? *S. fluitans* (E. M. Fries) E. M. Fries, Summa Vegetabilium Scand., 559, 1849 ≡ *S. simplex* var. *β fluitans* E. M. Fries, Flora Hallandica, pt. 1, 139. 1817. In 1849 Fries clearly bases *S. fluitans* on his "var. *fluitans*" from 1817. It is not possible to identify var. *fluitans* on the basis of its diagnosis in Flora Hallandica (pusillum, foliis decumbentibus natantibus, sequentis multo latioribus & brevioribus, basi 3-quertris). A type specimen must have been collected in Halland by 1817; no such specimen has been found even after a diligent search by Dr. Roland Moberg (UPS). In 1849 Fries wrote a new description of *S. fluitans* which bears almost no relation to his earlier diagnosis. This new description probably refers to *S. glomeratum* Laestadius; he also cites "Herb. normale XVI" (presumably a misprint for XV). From this evidence it is fairly clear that *S. fluitans* E. M. Fries and *S. glomeratum* Laestadius refer to the

same species. The earlier name is *S. fluitans* which is not only semantically unsuitable but has not been used for about a hundred years. Following the present interpretation of the Code of Botanical Nomenclature *S. fluitans* would not be accepted as a nomen rejiciendum and *S. glomeratum* would not be accepted as a nomen conservandum. We beg the botanical community to use *S. glomeratum* until a more sensible code of nomenclature exists.

= *S. glehnii* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4 (36): 34. December 1893. Type: USSR, "Sachalin, Sümpfe bei der Ansiedlung Tunai, 29 July, Glehn (holotype: LE, MHA or MW, not found by us but seen by Rother, 1910).

Robust but relatively small plants usually emergent with erect leaves and stems, rarely floating or submerged. Stolons up to at least 10 cm long, 1–1.5 mm diameter. Scale leaves ca. 10–18 mm long.

Basal leaves submerged or erect and distally emergent but rarely floating; submerged leaves thin and delicate, flat rarely more than 5 mm wide; emergent or floating leaves (20–)30–50(–60) cm long, (2–)4–9(–18) mm wide, in transverse section flat to very shallowly triangular distally, becoming shallowly triangular to deltate below with a winged abaxial keel, with 1 or below sometimes 2 layers of 6–8 gas chambers, not inflated at base.

Flowering stems (10–)30–40(–60) cm long, 2–4 mm diameter with 1–3(–4) elongated sterile nodes below the inflorescence, erect, robust, distinctly shorter than the leaves.

Inflorescence bracts usually straight and erect but remaining at an acute angle to the axis (Fig. 7 b), inflated at base with hyaline margins; lowermost bract (15–)20–30(–35) cm long, at least 3 times as long as the inflorescence; bract of the lowest male head usually brown and scale-like or sometimes absent. Inflorescence simple and compact.

Female heads (2–)3–5(–6), crowded, sessile or the lowest pedunculate and often remote, the rest usually supra-axillary, the lower ones often appearing above the next node or opposite the next bract; in fruit (10–)12–16(–20) mm diameter.

Male heads 1–2, more or less contiguous with the uppermost female head (Fig. 7 b).

Female flowers with perianth segments oblong to oblong-spathulate, with erose tips, clawed below, one third to half as long as the fruit, attached to a short pedicel; stigmas 0.6–0.8 mm long, lanceolate, obliquely attached to a 1.5–2.0 mm long style.

Male flowers with filaments 5–6 mm long; anthers narrowly oblong, 0.7–0.9 mm long.

Fruits narrowly fusiform, (3.0–)3.5–5.0(–6.0) mm long, 1.2–2.0 mm diameter, distinctly or indistinctly constricted near the centre, green-brown, shiny-surfaced, tapered below to an obconic base with a 0.5–2.0 mm long pedicel, tapered above to an acute summit with a conic based, 1.5–2.0 mm long beak; endocarp ovoid, 3.5–4.0 mm long, ca. 2 mm diameter, tapered below into a 2–3 mm long stalk, smooth with some very fine longitudinal furrows (Fig. 6 d).

Diagnostic features

Inflorescence simple; female heads crowded, the upper ones usually sessile, supra-axillary, often appearing above the next node or opposite the next bract; male heads 1–2, contiguous with uppermost female head; stigmas less than 0.8 mm long; mature fruit shiny with a straight beak; lowest inflorescence bract carinate to apex, at least 3 times as long as the inflorescence.

Distribution

S. glomeratum has a disjunct distribution (see Map 7 & 11). It is common and abundant in lowland and southern regions of Norway, Sweden and Finland, rarely reaching beyond 63°30'N (at Kuolajärvi it reaches to 67° N); eastwards it extends from White Russia to the Urals. It is apparently absent from Siberia and central Asia but reappears in the east where it grows in Manchuria, Liaoning, Jirin, N. Korea, Kamchatka and Japan (in Hokkaido and the mountains of central Honshu). There are some isolated records from eastern Tibet and Yunnan in S.W. China. There are also reports of *S. glomeratum* in N. America. We can confirm the record from Goose Bay, Labrador, Canada (20 July 1950, Gillet & Findlay 5390 - LU). The records of Lakela (1941) from Lake Country, Minnesota and Porsild (1950) from Dawson in the Yukon we have not been able to confirm and we doubt their validity (Map 11).

Ecology

S. glomeratum is essentially a species of shallow water in pools, small ponds, streams and ditches that have varying water levels and that may dry out in summer. The leaves are usually erect and mostly aerial; the aerial portions of the leaves are un-wettable on both surfaces. It is rarely found in large lakes but if so then it is confined to the edges in very shallow water. It shows a preference for mesotrophic neutral water, not being found in acid oligotrophic water and also avoiding calcium-rich waters. From the point of view of water quality it occupies a position between *S. natans* and *S. emersum*. In Scandinavia it is widespread east of the Kattegat but is by no means common only occasionally is it locally abundant; it is found growing with species such as: *Scirpus (Eleogiton) fluitans* L., *Eleocharis mamillata* (Lindb. f.) and *Ranunculus reptans* L. Among the northern species it is the first to flower and develop fruits; fruiting specimens have been collected as early as July.

Variation

The characteristic arrangement of the male and female heads is remarkably constant throughout the range of the species making *S. glomeratum* a highly distinctive member of the genus. Compared with other species of *Sparganium* it is not very variable. In Japan, however, some plants have narrow leaves which are flat (not keeled) and floating. These plants have been given varietal rank (var. *angustifolium*) by Graebner (1900); Hultén (1964) suggested that these plants belonged to another species. All floral characters are typical for *S. glomeratum* and without further biosystematic work we feel it unwise to give these plants formal taxonomic recognition.

Hybrids

Lid (1952) reports that *S. glomeratum* hybridizes with *S. angustifolium* and *S. emersum* in Norway. Yuzepchuk (1934) in Komarov reports *S. glomeratum* × *S. emersum* from Karelia-Lapland and Kamchatka. We have seen no convincing material of these hybrids, we doubt that the former exists but *S. glomeratum* × *S. emersum* may well exist but would be difficult to distinguish from *S. angustifolium* × *S. emersum*.

5. ***Sparganium gramineum* J. G. Georgi**, Bemerkungen einer Reise im Russischen Reich im Jahre 1772, 1: 232. 1775. Type: USSR, Buryatskaya ASSR, "in den Seen der oberen Angara, der Tanaga und dem Turtil", Georgi (we have failed to find any type

- specimens in LE and MW but they were known to Rothert, 1910) non Wallroth, Erster Beitrag Fl. Hercyn., 2: 297. 1840.
- = *S. lanceolatum* J. G. Georgi, Bemerkungen einer Reise im Russischen Reich im Jahre 1772, 1: 233. 1775. Type USSR, Buryatskaya ASSR, the same as *S. gramineum* (we have failed to find any type specimens in LE and MW but they were known to Rothert, 1910).
- = *S. friesii* Beurling, Bot. Not. 1854: 136. September–October 1854. Type: Sweden, Strömstad (lectotype: chosen here, “in lacu Strömvatnet, Jul. 1854, P. J. Beurling” S; isolectotype: S; paratype: “Ströms-än Julio 1954” manu Beurling, S.)

Slender plants with vegetative parts usually submerged or floating. Stolons not well developed (or rarely collected), short, ca. 3 mm diameter. Scale leaves up to 1 cm or more long, thin translucent. Basal leaves submerged below and floating distally, up to 120 cm or more long, (1–)2–3(–6) mm wide, not inflated at base, flat to plano-convex or semi-terete, with hyaline margins at base, without a distinct midrib; apices rounded.

Flowering stems 40–50(–200) cm or perhaps longer, 2–3 mm diameter, flexible, mostly submerged but emergent at tip; inflorescence simple or usually branched; main axis bearing (3–)5–7 female and 2–6 male heads; lowest lateral branch axillary bearing 1–3 female and (0–)1–2(–3) male heads, no other branches bear male heads; female and male heads separated by a (1–)2–4 cm long internode.

Inflorescence bracts at base inflated and with hyaline margins; lowermost bract up to 60 cm long, one to two times as long as the inflorescence (Fig. 7c); bracts of uppermost female heads and all male heads scale-like or absent.

Female heads always axillary, sessile or the lower shortly pedunculate, each remote and not touching, in fruit 10–20 mm diameter.

Male heads distinctly separated from female ones by a (1–)2–4 cm long internode, all more or less contiguous (Fig. 7c); axis remaining more or less straight after anthesis.

Female flowers with perianth segments oblong to spathulate, 2.5–3.5 mm long, erose at apex, translucent or with brown midrib, attached at base of pedicel, not obviously united below, at maturity about half as long as fruit; pedicel ca. 1 mm long; stigmas 0.6–0.7 mm long, ovate to triangular, obliquely attached.

Male flowers with filaments ca. 5 mm long; anthers (0.7–)0.8–0.9(–1.0) mm long.

Fruits ovoid, 2–3 mm long, 1.5–2 mm diameter, rarely constricted around middle, dark brown often with purple flecks, shiny-surfaced; tapering below to an obconic base with a 0.5–1.0 mm long pedicel; tapering above to conic-based beak; beak persistent, abruptly deflexed from base, 0.8–1.5 mm long; endocarp ovoid, ca. 2–2.5 mm long, somewhat constricted by ca. 6 shallow longitudinal furrows, bluntly conic at micropyle, elongated below into a persistent pedicel (Fig. 8a).

Diagnostic features

Leaves floating (apparently never erect), long and usually less than 5 mm wide; inflorescence simple or with a single branch; female heads axillary and remote, sessile or with the lowest shortly pedunculate; male heads 2–6, separated from the uppermost female by an internode at least 1 cm long; stigmas less than 0.8 mm long; fruit shiny with a short, abruptly deflexed, ca. 1 mm long beak.

Distribution

S. gramineum has a somewhat disjunct distribution (Map 8). In Scandinavia it is centred in S. and E. Sweden and S. Finland, mostly in low lying regions but reaching




Fig. 8. Perianth segments, fruits and endocarps of: a, *Sparganium gramineum*; b, *S. fluctuans*; c, *S. emersum* subsp. *emersum*; d, *S. emersum* subsp. *acaule* (scale bar: 1 mm).

346 m in Dalarna Jamtland and 292 m in Lappland. There are some isolated reports from E. Norway, extending to ca. 67° N.

In European Russia it grows from Leningrad and Novgorod north to Archangelsk and eastwards to Upper Volga (Moscow). It is absent from much of C. Russia but reappears in Siberia from Irkutskaya Oblast (around Lake Baikal) and Kamchatka.

Ecology

S. gramineum is more or less confined to oligotrophic lakes and is not or very rarely found in pools or ditches. It seems incapable of developing aerial leaves and may not be able to accommodate to changing water levels. The distal parts of the leaves float on the surface but they are apparently not so dorso-ventrally specialised as *S. angustifolium*. The submerged part of the leaf is often very thin and almost resembles a petiole. It is very gregarious and often develops very large populations; the floating leaves, which usually lie parallel, may cover the surface of the water. It is usually found in water from 40 cm to 150 cm deep with the consequence that it is rarely found near the banks. In depths of more than 1 m it often does not develop flowers and it is then virtually impossible to distinguish it from *S. angustifolium*. It has been recorded up to 3 m deep. It is unable to tolerate more than very small quantities of calcium, nitrogen and

phosphorous and is not found in brackish waters; it is a good indicator for oligotrophic water.

It is occasionally found growing with *S. angustifolium* with which it sometimes hybridizes. The hybrid apparently outcompetes the *S. gramineum* often leaving populations of *S. angustifolium* mixed with *S. angustifolium* × *S. gramineum*. *S. gramineum* also crosses with *S. emersum*; this hybrid is more tolerant of eutrophic conditions and rarely co-habits with pure *S. gramineum*.

Hybrid

5×3. *Sparganium gramineum* × *S. angustifolium*

From the morphology alone it is not possible to distinguish this hybrid from 5×7, *S. gramineum* × *S. emersum* (see p. 257). As *S. emersum* is a species of eutrophic waters it is to be expected that the latter hybrid is less common but the opposite seems to be the case. The hybrids tend to resemble *S. gramineum* but have longer stigmas and some supra-axillary heads. They are rather variable and characteristically have a somewhat irregular arrangement of the male heads (Fig. 9a & b). The hybrids are apparently fertile and the variability may result from segregation of self-fertilized hybrids or occasional backcrossing to either parent. These hybrids are relatively common in Scandinavia and regions where the distributions of the parent species overlap (see Map 9).

6. ***Sparganium fluctuans* (Engelmann ex Morong) B. L. Robinson, Rhodora, 7: 60. 1905 ≡ *S. androcladum* var. *fluctuans* Engelmann ex Morong, Bull. Torrey Bot. Club, 15: 78. 2 March 1888. Type: USA Pennsylvania, "ponds at the base of the White Mountains, *Oakes*" (holotype: MO n.v.).**

— *S. simplex* Hudson var. *fluitans* Engelmann in A. Gray, Manual Bot. Northern U.S., ed. 5, 481. 1867; nom. illeg., based on the type of *S. fluctuans*; non E. M. Fries, nec Döll, nec Godron & Grenier, nec Wirtgen. Type: USA, Pennsylvania, "ponds at the base of the White Mountains, *Oakes*" (holotype: MO n.v.).

Slender plants with vegetative parts usually submerged or floating. Stolons up to 20 cm or more long and 2–3 mm diameter.

Scale leaves up to 1 cm or more long, thin, translucent.

Basal leaves submerged below and floating distally, (20–)60–80(–100) cm long, (3–)4–8(–12) mm wide, not inflated at base, flat, thin, translucent, cross-reticulate abaxially, without a distinct midrib; apex rounded-acute.

Flowering stems (20–)60–100(–150) cm long, 1.5–3 mm diameter, mostly submerged, flexible, floating and emergent at tip; inflorescence usually branched with main axis bearing (0–)1(–2) female and (3–)4–6 male heads; lateral branches usually axillary (individual heads may be supra-axillary) bearing 1–2(–3) female and (1–)2–4 male heads (at distal end of inflorescence it is difficult to distinguish between main axis and branches, Fig. 7d); female and male heads separated by a distinct, (5–)10–20(–30) mm long internode.

Inflorescence bracts slightly inflated at base, usually with hyaline margins; lowermost bract (4–)6–18(–25) cm long, shorter than or sometimes slightly exceeding (up to 1.8 times) the inflorescence; bracts of uppermost female head and all male heads scale-like or absent.

Female heads axillary or supra-axillary, sessile, irregularly spaced along the axis or branches (Fig. 7d) at least some touching or almost touching, in fruit 15–23 mm diameter.




Fig. 9. Diagrammatic representation of inflorescences of: a, b, *Sparganium angustifolium* × *S. gramineum*; c, d, *S. emersum* × *S. gramineum*.

Male heads ca. 10–15 mm diameter, the lower ones distinct from each other, after anthesis leaving a zig-zag stalk, usually ebracteate.

Female flowers with perianth segments linear-oblong, 2–3.5 mm long, erose at apex, with dark brown tip and midrib, attached to the middle or below the middle of

the pedicel, not clearly united below, at maturity about half as long as the body of the fruit; pedicel ca. 1–3 mm long; stigmas (0.4–)0.7–0.8(–0.9) mm long, lanceolate-ovate, obliquely attached.

Male flowers with spathulate, ca. 3 mm long, dark-tipped perianth segments; filaments ca. 6 mm long; anthers (0.5–)0.6–0.7(–0.9) mm long.

Fruits ovoid to ovoid or fusiform, (2.7–)3.0–4.0(–5) mm long, ca. 2–2.5 mm diameter, usually constricted somewhat above the middle, dark brown, matt, dull-surfaced; tapering below to an obconic base with persistent 1.5–3 mm long pedicel; rounded above with a robust curved, 2–3.5 mm long beak; endocarp ovoid, ca. 3.5 mm long and 2.25 mm diameter, somewhat constricted by ca. 6 very shallow longitudinal furrows, flattened at the micropyle, elongated below into a persistent pedicel (Fig. 8 b).

Diagnostic features

Superficially resembling *S. gramineum* but generally more robust; inflorescence more branched with more heads on each branch; leaves usually more than 4 mm wide; beak curved, longer than 2 mm; stigmas usually more than 1 mm long; fruit usually more than 3 mm long with a pedicel exceeding 1.5 mm; anthers smaller, rarely exceeding 0.8 mm long. Easily distinguishable from the *S. americanum* group by the thin, translucent perianth segments.

Distribution

S. fluctuans is confined to boreal North America (Map 11) from Newfoundland westwards to northern Alberta and central British-Columbia extending southwards in the east to the northern counties of Pennsylvania.

Ecology

According to published accounts (Crow & Hellquist 1981, Brayshaw 1985) it is found in cold water ponds, lakes and slow rivers, often in relatively deep (1–2 m) water. It grows in oligotrophic usually somewhat acidic waters. It is reported to be uncommon in British Columbia and it is on the “rare and endangered plant list” for Connecticut.

Variation and hybrids

S. fluctuans is a distinct, rather invariable species that could be considered to be the New World vicariant of *S. gramineum*. Unlike *S. gramineum* it does not seem to hybridize with any other species.

7A. *Sparganium emersum* A. Rehmann, sensu stricto, Verhand. Naturforsch. Ver. Brünn, 10: 80. 1872 ≡ *S. simplex* proles *longissimum* subvar. *emersum* (Rehmann) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 286. 15 June 1897 ≡ *S. simplex* proles *longissimum* var. γ *emersum* (Rehmann) Graebner in Engler, Pflanzenreich, 2 (IV.10): 17. September 1900 ≡ *S. angustifolium* subsp. *emersum* (Rehmann) T.C. Brayshaw, Occasional Papers British Columbia Provincial Museum, 26: 115. 1985. Type: USSR, Ukraine, “Im Flussbette des Row bei Bar in Podolien”, 1868, A. Rehmann 154 (holotype: not found ?BRNM or BRNU; isotypes: B, K – paratype: Iter cilicico kurdicum, J. Kotschy 468: K).
= *S. erectum* L. var. β (*non ramosum*) L., Sp. Pl., 971. 1753. Type: Europe, specimen in Clifford’s Herbarium (BM).

- = *S. simplex* var. *gracilis* [sic] Meinshausen, Bull. Soc. Imp. Nat. Moscou, N.S., 3: 170. 1890 ≡ *S. simlex* var. *angustifolium* [sensu Morong] subvar. *gracile* [sic] (Meinshausen) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora, 1: 285. 15 June 1897 ≡ *S. simplex* var. γ *gracile* (Meinshausen) Graebner in Engler, Pflanzenreich, 2 (IV.10): 17. September 1900 [Graebner's citations loc. cit. are incorrect] ≡ *S. emersum* forma *gracile* (Meinshausen) Soó, Acta Bot. Acad. Sci. Hung., 17: 124. dated 1971, publ. 1972; non *S. natans* forma *gracilis* L. M. Neuman in C. J. Hartman & C. Hartman, Handbok Skand. Fl., ed. 12, 109. October 1889. Type: USSR, "Ingrien" "im nordlichen Torflande" (lectotype: chosen here, the specimen collected in July 1860 with "Teste Prof. Rothert" stamped on the label, LE).
- = *S. splendens* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4 (36): 32. December 1893 ≡ *S. simplex* var. *splendens* (Meinshausen) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora, 1: 285. 15 June 1897 ≡ *S. emersum* forma *splendens* (Meinshausen) Soó, Acta Bot. Acad. Sci. Hung., 17: 124. dated 1971, publ. 1972. Type: USSR, "nur in Süd Gebiete Ingriens gefunden" (lectotype: chosen here, sheet bearing label "*Sparganium splendens* m. n.sp" and another label with handdrawn crown and "n.sp exellentissima" written on it, LE). Some iso- or syntypes may represent the hybrid *S. angustifolium* × *S. emersum*.
- = *S. simile* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4 (36): 34. December 1893 ≡ *S. simplex* var. *splendens* subvar. *simile* (Meinshausen) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 285. 15 June 1897 ≡ *S. simplex* var. ϱ *simile* (Meinshausen) Graebner in Engler, Pflanzenreich, 2 (IV.10): 17. September 1900 ≡ *S. emersum* forma *simile* (Meinshausen) Soó, Acta Bot. Acad. Sci. Hung., 17: 124. dated 1971, publ. 1972. Type: USSR, "monte Calmytolohey in deserto Siungorico, 1771, Falk" (holotype not found in LINN, LE, MW).
- = *S. subvaginatum* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg, N.S., 4 (36): 34. December 1893, pro parte ≡ *S. simplex* var. *angustifolium* subvar. *subvaginatum* (Meinshausen) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 285. 15 June 1897 ≡ *S. simplex* var. δ *subvaginatum* (Meinshausen) Soó, Acta Bot. Acad. Sci. Hung., 17 (1–2): 124. dated 1971, publ. 1972. Type: Finland, Archipelago Aboensi-Krånskor, pr. Kelo, Ruprecht; Ins. Sitcha, Mertens; Rocky Mountains, reg. alpina, C. C. Parry (lectotype to be chosen).
- = *S. diversifolium* Graebner, Schriften Naturf. Ges. Danzig., N.F., 9 (1): 335, 1896, pro parte. Type: Poland "Bielawa-Bruch ... Slawoschin ... Lübtower See, leg. A. Triechel ... Wobrow, leg. Graebner" (holotype: ?destroyed in B). In a letter to Fernald (cited in Rhodora, 24: 29. 1922) Rothert wrote "[Graebner's name [*S. diversifolium*] comprises chiefly *S. simplex* × *minimum* [*S. emersum* × *natans*] and slender forms of *S. simplex*; besides, I have seen specimens of *S. affine* [*S. angustifolium*], *S. minimum*, *S. glomeratum* and *S. affine* × *minimum* determined by G. himself as *S. diversifolium*, not a single one of all these fitting his description." We have also seen different taxa determined as *S. diversifolium* by Graebner in B.
- = *S. chlorocarpum* Rydberg, North American Flora, 17: 8. 30 June 1909 ≡ *S. angustifolium* subsp. *emersum* var. *chlorocarpum* (Rydberg) T. C. Brayshaw, Occasional Papers British Columbia Provincial Museum, 26: 115. 1985. Type: USA, Iowa, Emmet County, "rare in marshes", September 1898, Alta Cratty s.n. (holotype: NY).
- = *S. diversifolium* proles *wirtgeniorum* Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 290. 15 June 1897 ≡ *S. wirtgeniorum* (Graebner) Rouy, Fl. France, 13: 339. 1912. Type: Germany, Rheinland, "Laacher See, Rodder Maar, Mülheim b. Köln, Viersen" Wirtgen (holotype: ? destroyed in B).

- = *S. simplex* proles *longissimum* subvar. *inundatum* Schur ex Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 286. 15 June 1897 = *S. simplex* proles *longissimum* var. β *inundatum* (Schur ex Graebner) Graebner in Engler, Pflanzenreich, 2 (IV.10): 17. September 1900. Type: Austria, "im Prater bei Wien, Schur" (holotype: ? destroyed in B).
- *S. simplex* Hudson, Flora Anglica, ed. 2, 2: 401. 1778 = *S. erectum* var. α *simplex* (Hudson) Laestadius, Bihang Wikström's Årberättelse, 1850: 1. 1853 or 1854 = *S. emersum* subsp. *simplex* (Hudson) Soó, Acta Bot. Acad. Sci. Hung., 17: 124. dated 1971, publ. 1972; nom. illeg., based on *S. natans* L., see Cook (1985).
 - *S. simplex* var. *longissimum* E. M. Fries, Bot. Not., 1868: 71. 1868 = *S. simplex* proles *longissimum* (E. M. Fries) Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora, 1: 285. 15 June 1897 = *S. longissimum* (E. M. Fries) Fritsch, Exkursionsfl. Österreich, ed. 2, 29. 1909; nom. illeg., based on *S. erectum* var. β *boreale* Laestadius.
 - *S. simplex* var. *angustifolium* Beckmann, Abhandl. Natforsch. Ver. Bremen, 10: 505. 1889; nom. illeg., combination made in 1867 by Engelmann for "*angustifolium*" Michaux.
 - *S. emersum* forma *natans* (Glück ex) Soó, Acta Bot. Acad. Sci. Hung., 17: 124. dated 1971, publ. 1972; nom. illeg., Glück's "Formen" represented phenotypically induced states.
 - *S. emersum* forma *submersum* (Glück ex) Soó, Acta Bot. Acad. Sci. Hung., 17: 124. dated 1971, publ. 1972; nom. illeg., Glück's "Formen" represented phenotypically induced states.

Robust to slender, submerged, floating or erect herbs.

Stolons up to ca. 50 cm long, (1-)2-3(-4) mm diameter. Leaves of fertile plants usually erect and partially emergent, (10-)20-50(-80) cm long and (1.5-)4-10(-12) mm wide, usually distinctly carinate from base to apex, with 1-3(-5) layers of up to 12 gas chambers.

Plants in deep or swiftly flowing water usually remain sterile; leaves of sterile plants submerged or floating, flat to somewhat keeled, up to 220 cm long and 18 mm wide.

Flowering stems simple, erect, 20-60(-80) cm long or when floating up to ca. 180 cm long, with 1-3 sterile nodes between the corm and the inflorescence.

Inflorescence bracts usually erect and carinate at base with hyaline margin and sometimes somewhat inflated; lowermost bract up to ca. 25 cm long, 1-2(-4) times as long as the inflorescence; bract of upper female head usually not exceeding the inflorescence (Fig. 10 a & b).

Female heads (1-)3-4(-6), the lower usually pedunculate and axillary (peduncles 0-4(-8) cm long), the upper sessile and some usually supra-axillary; in fruit 1.6-2.5(-3.5) cm diameter, appearing white to yellowish-green at anthesis.

Male heads (3-)4-7(-10), remote and distinct at anthesis, separated from the uppermost female head by a 3-20(-40) mm long internode; before anthesis pale yellowish-green.

Female flowers with perianth segments spathulate, translucent, with erose tips, scarcely clawed below (Fig. 2a), one-half to two-thirds as long as the fruit, united at base usually attached to pedicel (pedicel 1.0-2(-3) mm long); stigmas (1.0-)1.5-2(-2.5) mm long, obliquely attached to a 2-3 mm long style.

Male flowers with filaments 5-7 mm long; anthers 1-1.5(-2) mm long.

Fruits fusiform, sometimes slightly constricted around the middle, 3.5-5.5 mm long, ca. 1.8-2.5 mm wide, brown, shiny, tapering above to a 2.0-4.5(-6) mm long beak, tapering below to an obconic base with an up to 4 mm long (by shrinkage of the fruit)




Fig. 10. Diagrammatic representation of inflorescences of: a, *Sparganium emersum* subsp. *emersum* (typical European inflorescence); b, *S. emersum* subsp. *emersum* (typical N. American inflorescence); c, *S. emersum* subsp. *acaule*; d, *S. angustifolium* × *S. emersum* subsp. *emersum*.

stalk; endocarp ovoid, 2.5–3.5 mm long, ca. 1.8–2.5 mm diameter, tapered below to a 1–4 mm long stalk, smooth with some fine longitudinal ridges (Fig. 8 c).

Diagnostic features

Inflorescence simple; male heads 3 or more, remote and distinct at anthesis; stigmas more than 1 mm long; anthers 1–2 mm long; perianth segments translucent (heads appearing white to greenish-yellow at anthesis); upper female heads usually sessile and supra-axillary; leaves and lower bracts of flowering plants usually erect and distinctly carinate to apex. It may superficially resemble species of the *S. americanum* group; the presence of thin, translucent perianth segments is the distinguishing feature, see also Table 1 in part 2 (subgenus *Sparganium*).

Distribution

S. emersum has a holoarctic distribution with the main area in the temperate zone, Maps 12 and 13. It is found on lower ground throughout Europe from about the arctic circle to the submediterranean zone. In Asia it extends from the arctic circle south to about 40° N (growing between 1900 and 2300 m in Anatolia), eastward to Japan; there are isolated occurrences south of 40° N in high land in the Gilgit Wazerat district, around Nanga Parbat, Northern Area of Kashmir (3000 m) and Helan Shan (China).

In America it extends from east to west, from south of the arctic circle to about 40° N; in the southern parts of its range it is usually found at high levels, reaching 2900 m in Colorado and 2500 m in Wyoming.

The record from Sumatera is incorrect; it refers to *S. fallax*.

Ecology

We have experience of *S. emersum* in the field in central, west and southern Europe and have seen it growing in western N. America. It is found in a wide variety of aquatic habitats in still and flowing water but usually grows near the bank, particularly in somewhat disturbed or unstable places. Stolons are well-developed but *S. emersum* very rarely becomes dominant and is usually found in relatively small stands (up to about 12 m²) or isolated clumps of shoots. Although it can tolerate periods of drought it is essentially an aquatic species that is partly or totally submerged most of the time.

S. emersum is relatively deep-rooted (15 cm or more deep) and is very tolerant to disturbance. For example, it is frequent in *Phragmites* or other reed-swamp communities in clearings or channels that are artificially maintained (for boating, fishing, swimming etc.); it also grows in ricefields in Europe without however, becoming a serious weed. It is mostly found in places where it is subjected to rapid changes in water level or irregular flooding.

Plants are not easily dislodged and it can stand severe spates. The leaves are somewhat frail and are frequently damaged following floods or disturbance but they seem to be quickly replaced. In a stream in Denmark, Waggers Neilson et al. (1985) found the lifetime of individual leaves was only 31–39 days and that the mortality rate constituted more than 50% of the production rate before maximum biomass was attained. These findings elegantly demonstrate that standing crop cannot be used as a measure for annual production.

It is usually found in shallow water (20–80 cm deep) but has been recorded in water up to 2.6 m deep in Finland. The leaves of *S. emersum* rarely grow more than 2 m long and they are usually much shorter so that plants in deep or in swiftly-flowing water remain submerged. Water quality does not seem to be very important; although it is most common in mesotrophic to semieutrophic conditions, it may occasionally be found in almost oligotrophic or strongly eutrophic water but is not tolerant to brackish water and is rarely found in salinities exceeding 0.2%. The substrate preferences are rather more restricted. It grows best in deep clay or fine, sandy soils. It is a characteristic species in the silted parts of small rivers and streams. In meandering rivers one tends to find it at almost each curve. It is also found along the banks of ponds, lakes, canals and drainage ditches. In Europe, it is ecologically rather like *Sagittaria sagittifolia* and both species frequently grow in an association named Sparganieto-Sagittarietum.

Sparganium emersum shows a rather wide ecological spectrum but is a poor competitor which probably accounts for its inability to become dominant. Its performance is however not alike in all habitats and it is frequently found in a nonflowering state. For flowering it seems to need shallow water (less than 50 cm deep), a lot of light and little competition. Its poor ability to compete is compensated by its ability to withstand disturbance. Nevertheless, it does not possess the high growth rate that one expects from a ruderal (Grime 1979). As it grows in and is active in collecting and consolidating silt it is essentially a species in the middle of the aquatic hydrosere whereby aquatic habitats are converted into terrestrial ones.

Variation

S. emersum is a polymorphic species showing variation in the following characters: leaf-width, number and position of male and female heads, stigma length and anther length. In the past this has led to differing opinions regarding the taxonomic boundaries of the species and particularly in distinguishing it from *S. angustifolium*. The situation is complicated by the occurrence of hybrids (see p. 255).

Packer (1983) has reviewed these taxonomic failings. Comparing two recent keys to the species (Reveal, in Cronquist et al. 1977, and Cook, in Tutin et al. 1980) he found some surprising differences. Reveal describes *S. angustifolium* as having 2–5 male heads and in the accompanying illustration these are clearly depicted as being remote. The account also describes the leaves of both species as flat or rounded (but considers those of *S. emersum* as being weakly keeled underneath). Neither of these assertions are correct. It is unfortunate that Reveal's account should confuse these two attributes since they are of major importance in understanding the distinction between the species. The leaves of *S. angustifolium* are flat or abaxially convex but those of *S. emersum* are clearly triangular, at least, at the base. The most striking single attribute of *S. angustifolium* is that the male heads are less than four in number and contiguous (giving the appearance of a single elongated head); this is never the case in *S. emersum*. While Packer (1983) is correct in recognizing these difficulties, his decision to combine *S. emersum* (and *S. chlorocarpum*) under the single specific name *S. angustifolium* only makes matters worse.

The position of the female heads of *S. emersum* ranges from axillary and sessile to supra-axillary and pedunculate. There appears to be little geographic consistency in these attributes with the exception of subsp. *acaule* (see p. 257) which is confined to the northeast of North America; it has female heads that are typically crowded, supra-axillary and sessile. Generally, large and robust plants of *S. emersum* have the lower female heads pedunculate while small depauperate plants have fewer and sessile heads.

The position and number of the male heads also varies; while typically distinct at anthesis, the immature heads can be observed crowded at the distal end of the inflo-




Fig. 11. Map representing the length of the stigmas of *Sparganium emersum* in Eurasia.

rescence. Variation in stigma length and anther length is strongly related to geography (see Fig. 11). Plants from west and central Europe possess relatively small stigmas (1.0–1.5 mm long) and anthers (1.0–1.3 mm long). Specimens from Scandinavia, eastern Europe, Turkey, Afghanistan and Siberia have longer stigmas (1.5–2.0 mm long) and anthers (1.3–1.6 mm long). The variation pattern is clinal from east to west.

Specimens of *S. emersum* from northwest North America often show characteristics of *S. angustifolium*. Particularly confusing is the number and arrangement of male heads; often there are three or four heads, the lowest is remote but often the upper are somewhat contiguous (Fig. 10 b). Also the lowest bracts are often much longer than the inflorescence. The habit, arrangement of female heads and the dimensions of the flowers are all typically *S. emersum*. Populations appear to be constant in form and there is no evidence of segregation. It is possible that these plants are "stabilized" hybrids or backcrosses involving *S. angustifolium* but we agree with Rothert (determinations on herbarium sheets) that these plants are nevertheless assignable to *S. emersum*.

Several authors, including Casper & Krausch (1980) are convinced that the plants of deep or flowing water with long, band-like leaves are taxonomically distinct from the erect, partly emergent plants. Although the plants in extreme states do look very different, there is no convincing evidence to suggest that these states are genetically distinct. It is at times tempting to give formal taxonomic recognition to extreme phenotypes but is it not currently in accordance with taxonomic practice.

Hybrids

The following hybrids have been reported in nature (none have been raised after crossing experiments):

1×7. *Sparganium hyperboreum* × *S. emersum*. These species are largely allopatric. However, in Scandinavia they sometimes grow near together and Lid (1952) has recorded hybrids. From the available herbarium material we are not convinced that this hybrid exists.

2×7. *Sparganium natans* × *S. emersum* (often but incorrectly named *S. wirtgeniorum* (Graebner) Rouy). Many workers including Rothert (1910, 1913) are convinced that this hybrid exists. Both species often grow near together so one might expect hybrids where they are sympatric. We have seen no herbarium material that we consider to belong to this hybrid, also in the field we have never found hybrid plants in regions where both species grow together. We therefore doubt that this hybrid exists. In any case, on morphological grounds alone it may be difficult to distinguish it from some segregants of *S. emersum* × *S. angustifolium*, *S. emersum* × *S. glomeratum* or *S. emersum* × *S. gramineum*.

3×7. *Sparganium angustifolium* × *S. emersum*.

= *S. diversifolium* Graebner, pro parte, Schriften Naturforsch. Ges. Danzig, N.F., 9 (1): 335. 1896, pro parte. Type: Poland, "Lübtower See", "Bielawa-Bruch" (holotype: destroyed in B).

– *S. boreale* Beurling, Öfversigt Kongl. Vetenskaps-Akad. Förhandl. (Stockholm), 9 (8): 192, 13 October 1852, nomen nudum = *S. erectum* var. β *boreale* (Beurling) Laestadius, Bihang till Wikström's Årberättelse, 1850: 1. ?1853, nomen illeg., based on *S. natans* L. (authentic material of *S. boreale* is in S; see Cook 1985).

– *S. × zetlandicum* G. C. Druce, British Plant List, ed. 2, 115. 1928, nomen nudum.

– (*S. borderi* [sic] W. O. Focke is often considered to be a hybrid but the type material is clearly terrestrially grown *S. angustifolium*).




Fig. 12. A pictorialized scatter diagramme showing the relationship between anther length (vertical axis) and stigma length (horizontal axis) in *Sparganium angustifolium* (closed circle), *S. emersum* (diamond) and *S. angustifolium* \times *S. emersum* (circle with cross).

Hybrids are intermediate between the parents in vegetative and floral characters. The F_1 is highly fertile. Segregants and backcrosses are frequent and an almost continuous variation pattern between the parents is found. The F_1 has the inflated leaf-bases of *S. angustifolium* (even when growing terrestrially) but the remote male heads of *S. emersum* (Fig. 10d). Stigma length and anther size are intermediate (Fig. 12).

S. emersum is mostly found in eutrophic conditions while *S. angustifolium* is confined to more or less oligotrophic waters. Most of the range of *S. angustifolium* lies within that of *S. emersum*. In northern Europe hybrids are common and usually grow with one or both parent species. In northwestern North America there is evidence that some *S. angustifolium* characters have become incorporated into *S. emersum* and that "pure" *S. emersum* (as it is realized in Europe) has been replaced by a relatively stable hybrid (Fig. 10b).

In western Scotland eutrophic and oligotrophic pools are often found near together; in Stoer, Wester Ross (now Highland) a hybrid population was seen (by C.D.K.C.) growing on a shingle beach, a habitat not exploited by either parent.

4 \times 7. *Sparganium glomeratum* \times *S. emersum*. In 1910 Rothert omitted this hybrid from his account of *Sparganium* hybrids but in 1916 he lists it without any description. Lid (1952) reports it from southern Scandinavia. We have seen no convincing material but feel that it may well exist but would be very difficult to distinguish from *S. angustifolium* \times *S. emersum* (see above).

Table 3. Comparison of *S. emersum*, *S. gramineum* and their hybrid.

Character	<i>S. emersum</i>	Hybrid	<i>S. gramineum</i>
Inflorescence	Simple	Usually branched	Usually branched
Lowest ♀ bract/inflorescence ratio	1.0–4.0	1.3–3.4	1.0–2.0
Position of upper female heads	Supra-axillary	Supra-axillary	Axillary
Male head number	3–10	3–7	2–6
Male heads at anthesis	Distinct	Contiguous	Contiguous
Stigma length (mm)	More than 1.0	0.6–1.0	0.6–0.7
Beak	Straight	Straight or curved	Deflexed

5 × 7. *Sparganium gramineum* × *S. emersum*.

= *S. speirocephalum* L. M. Neuman in Hartman, C. J. & Hartman, C., Handbok Skand. Flora, ed. 12, 109. October 1889, pro parte. Type: Sweden (lectotype to be chosen from material in UPS; isotype B).
– *S. longifolium* Turczaninow Bull. Soc. Imp. Nat. Moscou, 1: 103. 1838. nomen illeg., nomen nudum. “Type”: USSR, Buryatskaya, ASSR, Daurica, “Ad fl. Angaram superiorum, 1834, Turzmannoff” (K).

This hybrid is common in regions where both parents grow together (Map 10). The hybrid is much more tolerant of eutrophic conditions than *S. gramineum* and may occupy areas where *S. gramineum* is absent or in habitats where *S. gramineum* has become extinct due to increasing eutrophication. The hybrid is fully fertile and while some segregation or back-crossing may be evident it is usually distinct (see Table 3 and Fig. 9 c, d).

This hybrid is often difficult to distinguish from *S. gramineum* × *S. angustifolium* (Fig. 9 a, b). The latter hybrid usually has 2–3 contiguous male heads, inflated bases to the inflorescence bracts, a simple inflorescence with some supra-axillary female heads and it grows in oligotrophic water.

7B. ***Sparganium emersum* subsp. *acaule*** (Beeby ex Macoun) C. D. K. Cook & M. S. Nicholls, comb nov. ≡ *S. simplex* var. *acaule* Beeby ex Macoun, Geological & Natural History Survey of Canada, Cat. Canadian Plants, Part 5 (Acrogens): 367. 1890 ≡ *S. diversifolium* var. *acaule* (Beeby ex Macoun) Fernald & Eames, Rhodora, 9: 88. May 1907 ≡ *S. acaule* (Beeby ex Macoun) Rydberg, North American Flora, 17: 8. 30 June 1909 ≡ *S. chlorocarpum* var. *acaule* (Beeby ex Macoun) Fernald, Rhodora, 24: 29. February 1922 ≡ *S. chlorocarpum* forma *acaule* (Beeby ex Macoun) E. G. Voss, Rhodora, 68: 436. October–December 1966. Type: Canada, “Quite common in ponds and wet spots by the road-side in many parts of Prince Edward Island, especially at Lake Verde, Brackley Point and Winter River”, 1888, Macoun (lectotype to be chosen at CAN).

= *S. diversifolium* proles γ *nanum* Graebner in Engler, Pflanzenreich, 2 (IV. 70): 21. September 1900. Type: North America: “Neuengland: White Mountains (Tucker- man jun. !); Terra Nova (*Lapyliae*!)” holo- or lectotype destroyed in B.

Like *S. emersum* sensu stricto but usually smaller and less robust; leaves 3–7 mm wide; basal leaves and lower inflorescence bracts strongly erect and conspicuously longer than the flowering stems (Fig. 10c); female heads supra-axillary, sessile and crowded (the lowermost sometimes remote and pedunculate); anthers 0.8–1.0 mm

long; stigmas 0.8–1.5(–1.7) mm long; fruits 3–4 mm long, ca. 1.5 mm wide, light brown, constricted around the middle; beak 3–4 mm long, about equal to the fruit body in length (Fig. 8d).

Distribution

It is confined to eastern North America (Map 13), extending from Newfoundland southwards to Virginia and reaching westwards south of the Great Lakes to Minnesota and Iowa. It does not totally replace *S. emersum* subsp. *emersum* in this region although it is by far the most frequent of the two.

Notes

In its characteristic form, *S. emersum* subsp. *acaule* is easily distinguished on account of its erect habit and constricted female heads. Nevertheless, detailed studies suggest that a range of intermediates exist between it and subsp. *emersum* and that since the major differences are qualitative, it is often difficult to distinguish the two with confidence. In view of this and its limited distribution we have recognized it at the rank of subspecies. Its ecology is probably like *S. emersum* subsp. *emersum* but from examination of herbarium material it would seem to grow better in very shallow water.

Résumé

Révision taxonomique complète du genre *Sparganium* (Sparganiaceae) avec clé, et descriptions complètes de chaque espèce y compris les diagnoses, synonymes avec typifications, cartes de distribution, variabilité et illustrations qui elles-mêmes contiennent des informations sur les fossiles, l'anatomie, la morphologie, les chromosomes, la biologie florale, la propogation, l'écologie, les parasites et animaux prédateurs, des aspects appliqués ainsi que les hybrides. Quatorze espèces et six sous-espèces sont reconnues. Dans la partie 1, sous-genre *Xanthosparganium*, 7 espèces sont présentées. Un seul nouveau nom est employé: *Sparganium emersum* subsp. *acaule* (Beeby ex Macoun) C. D. K. Cook & M. S. Nicholls.

References


- Abuziarova R. J. 1970. Spore-pollen complexes in the south of the Altai, locality Bukhtama (in Russian). Botaničeskij Žurnal Akad. Nauk SSSR. (Leningrad) 55: 1121–1126.
- Arohonka S. 1982. Kromosomilukumääritysä Nauvon Seilin saaren putkilokasveista Turun Yliopiston Biologian Laitoksen Julkaisuja, 3: 1–12; n.v. cited by Uotila P. & Pellinen K., 1985. Acta Bot. Fennica 130: 1–37.
- Benum P. 1958. The Flora of Troms Fylke. Tromsø Museums Skrifter 6: 82–84.
- Berry E. W. 1924. A. *Sparganium* from the Middle Eocene of Wyoming. Bot. Gaz. 78: 342–348.
- Bijok K. & Mlynek T. 1965. Badania cytologiczne nad trzema gatunkami rodzaju *Sparganium* L. Zesz. Nauk WSR. w Olsztynie 19: 205–211.
- Bijok K. & Adamkiewicz E. 1971. Badania kariologiczne nad dwoma gatunkami rodzaju *Sparganium* [sic]. Acta Soc. Bot. Poloniae 40: 143–148.
- Brayshaw T. C. 1985. Pondweeds and Bur-reeds, and their relatives, of British Columbia. Occasional Papers of the British Columbia Provincial Museum 26: 1–167.

- Casper S. J. & Krausch H.-D. 1980. Pteridophyta und Anthophyta. 1. Teil: Lycopodiaceae bis Orchidaceae. Süßwasserflora von Mitteleuropa, ed. Ettl H., Gerloff J. & Heynig H., 23: 76–91. VEB Gustav Fischer Verlag, Jena.
- Chen Yao-Dong 1981. A study on Chinese *Sparganium* (in Chinese). Acta Phytotaxonomica Sinica 19: 41–57.
- Cook C. D. K. 1961a. *Sparganium* in Britain. Watsonia 5: 1–10.
- Cook C. D. K. 1961b. Die bayerischen *Sparganium*-Arten Ber. Bayer. Bot. Ges. 34: 7–10.
- Cook C. D. K. 1962. Biological Flora of the British Isles, No. 82: *Sparganium erectum* L. J. Ecology 50: 247–255.
- Cook C. D. K. 1963. *Sparganium* nomenclature. Proc. Bot. Soc. British Isles 5: 124.
- Cook C. D. K. 1985. *Sparganium*: some old names and their types. Bot. Jahrb. Syst. 107: 269–276.
- Cronquist A., Holmgren N., Reveal J. & Holmgren P. 1977. Intermountain Flora 6: 465–468. Columbia Univ. Press, New York.
- Crow G. E. & Hellquist C. B. 1981. Aquatic vascular plants of New England: Part 2. Typhaceae and Sparganiaceae. New Hampshire Agric. Exper. Station Bull. 517: 1–21.
- Dietz S 1887. Über die Entwicklung der Blüte und Frucht von *Sparganium* Tourn. und *Typha* Tourn. Bibliotheca Botanica 5: 1–56.
- Dorofeev P. I. 1979. On taxonomy of Tertiary *Sparganium* (in Russian). Sovetskaya palaeokar-pologya. Publishing House Nauka, Moscow, 53–75.
- Duke J. A. & Ayensu E. S. 1985. Medicinal plants of China. Vol. 2, p. 614. Reference Publishing Inc., Algonac, Michigan.
- Feráková V., in Májovský J. 1974. Index of chromosome numbers of Slovakian flora. Acta Fac. Rerum Nat. Univ. Comenianae, Bot., 23: 1–23.
- Fernald M. L. & Eames A. J. 1907. Preliminary lists of New England plants. – XX. Sparganiaceae. Rhodora 8 (85): 86–90.
- Gadella T. W. J. & Kliphuis E. 1973. Chromosome numbers of flowering plants of the Netherlands. Proc. Koninkl. Nederl. Akad. Wetenschappen, ser. C, 76: 303–311. (repr.: Mededel. Bot. Mus. Herb. Rijksuniv. Utrecht, No. 392: 303–311).
- Gervais C. 1981. Liste annotée de nombres chromosomiques de la flore vasculaire du nord-est de l'Amérique. II. Naturaliste Canad. 108: 143–152.
- Glück H. 1938. Über das Vorkommen des arktischen *Sparganium hyperboreum* in der zentraleuropäischen Alpenkette. Bot. Jahrb. 69: 220–251.
- Graebner P. O. P. 1900. Sparganiaceae, in Engler H. G. A., Das Pflanzenreich 2 (IV.10): 10–24.
- Grime J. P. 1979. Plant strategies and vegetational processes. John Wiley & Sons, Chichester, New York, Brisbane, Toronto, pp. 1–222.
- Guppy H. B. 1906. Observations of a naturalist in the pacific between 1896 and 1899. Vol. 2, Plant-Dispersal. Mac Millan, London, pp. 627.
- Hagerup O. 1941. Nordiske Kromosom-Tal. I. Bot. Tidsskr. 45: 385–403.
- Harada I. 1949. Chromosome numbers in *Pandanus*, *Sparganium* and *Typha*. Cytologia 14: 214–218.
- Harms V. L. 1973. Taxonomic studies of North American *Sparganium*, I *Sparganium hyperboreum* and *S. minimum*. Canad. J. Bot. 51: 1629–1641.
- Hegelmaier F. 1874. Zur Entwicklungsgeschichte monokotyledoner Keime nebst Bemerkungen über die Bildung der Samendeckel. Bot. Zeitung 32 (29): 631–640.
- Hultén E. 1964. The circumpolar plants. Kungl. Svenska Vetenskapsakad. Handl. F.S. 8 (5): 1–280.
- Jørgensen C. A., Sørensen T. H. & Westergaard M. 1958. The flowering plants of Greenland. A taxonomical and cytological survey. Kongl. Danske Videnskab. Selskab. Biol. Skrift. 9 (4): 1–172.
- Kaul R. B. 1972. Adaptive architecture in emergent and floating *Sparganium*. Amer. J. Bot. 59: 270–278.
- Kaul R. B. 1973. Development of foliar diaphragms in *Sparganium eurycarpum*. Amer. J. Bot. 60: 944–949.
- Kaul R. B. 1976. Anatomical observations on floating leaves. Aquatic Bot. 2: 270–278.


- Kirchner O., Löw E. & Schröter C. 1908. Lebensgeschichte der Blütenpflanzen Mitteleuropas 1 (1): 374–394. Eugen Ulmer, Stuttgart.
- Kožuharov S. & Kuzmanov B. 1964. Chromosome numbers of four Bulgarian plants. Comptes Rendus Acad. Bulgare Sci. 17 (5): 491–494.
- Lakela O. 1941. *Sparganium glomeratum* in Minnesota. Rhodora 43: 83–85.
- Larsen K. 1965. IOPB Chromosome number reports. III. Taxon 14: 50–57.
- Leereveld H. 1984. Anthecological relations between reputedly anemophilous flowers and syrphid flies. VI. Aspects of the anthecology of Cyperaceae and *Sparganium erectum* L. Acta Bot. Neerl. 33: 475–482.
- Lid J. 1952. Norsk Flora. Det Norske Samlaget, Oslo. pp. 1–771.
- Löve Á. & Löve D. 1942. Chromosome numbers of Scandinavian plant species. Bot. Not. 1942: 19–59.
- Löve Á. & Löve D. 1948. Chromosome numbers of northern plant species. Atvinnudeild Háskóla Íslends Rit Landbúnaðar-deilar, B-flokkur 3: 9–131.
- Löve Á. & Löve D. 1956. Cytotaxonomical conspectus of the Icelandic flora. Acta Horti Gothburgensis 20 (4): 65–291.
- Löve Á. & Löve D. 1975. IOPB chromosome number reports XLVIII. Taxon 24: 367–372.
- Löve Á. & Löve D. 1981. Chromosome number reports LXXII. Taxon 30: 694–708.
- Löve Á. & Ritchie J. C. 1966. Chromosome numbers from central Canada. Canad. J. Bot. 44: 429–439.
- Mai D. H. 1985. Entwicklung der Wasser- und Sumpfpflanzen-Gesellschaften Europas von der Kreide bis ins Quartär. Flora 176: 449–511.
- Martin A. C. & Uhler F. M. 1939. Food of game ducks in the United States and Canada. Techn. Bull. US. Dep. Agric. 634: 1–157.
- Mehra P. N. & Pandita T. K. 1979. IOPB chromosome number reports LXIV. Taxon 28: 391–408.
- Muller J. 1984. Significance of fossil pollen for angiosperm history. Annals Missouri Bot. Garden 71: 419–443.
- Müller-Doblies Ute 1969. Über den Blütenstand und Blüten sowie zur Embryologie von *Sparganium*. Bot. Jahrb. Syst. 89: 359–450.
- Müller-Doblies Dietrich 1970. Über die Verwandtschaft von *Typha* und *Sparganium* im Infloreszenz- und Blütenbau. Bot. Jahrb. Syst. 89: 451–562.
- Müller-Doblies Ute & Müller-Doblies Dietrich 1977. *Sparganium* in Hegi G., Illustr. Flora Mitteleuropa, ed. 3, II, 1 (4): 275–317.
- Murin A. in Májovský J. 1978a. Index of chromosome numbers of Slovakian flora (Part 6) Acta Fac. Rerum Nat. Univ. Comenianae, Bot., 26: 1–42.
- Murin A. in Májovský J. 1978b. IOPB chromosome number reports LXI. Taxon 27: 375–392.
- Packer J. G. 1983. Flora of Alberta: *Sparganium angustifolium* and *Erigeron trifidus*. Canad. J. Bot. 61: 359–366.
- Pinkess L. H. 1980. The possibility of pollination of *Sparganium erectum* by insects. Proc. Birmingham Nat. Hist. Soc. 24: 101–102.
- Porsild A. E. 1951. Botany of Southeastern Yukon adjacent to the Canol road. Nat. Mus. Canad Bull. 121: 76.
- Praeger R. L. 1913. On the buoyancy of the seeds of some Britanic plants. Scient. Proc. Royal Dublin Soc. 14: 13–62.
- Reveal J. L. 1970. *Sparganium simplex* Huds., a superfluous name. Taxon 19: 796–797.
- Rothert W. 1910. Übersicht der Sparganien des Russischen Reiches (zugleich Europa's). Acta Horti Bot. Univ. Imper. Jurjevensis 11: 11–32.
- Rothert W. 1913. In Fedtschenko B. A., Flora Aziatskoi Rossii 1: 17–37, St. Petersburg.
- Saccardo F. 1895. Ricerche sull'anatoma della Typhaceae. Malpighia 9: 3–30.
- Scheerer H. 1940. Chromosomenzahlen aus der schleswig-holsteinischen Flora. I. Planta 29: 636–642.
- Sokolovskaya A. P. 1963. Geographic distribution of polyploid plants (observations from the Kamchatka Peninsula), in Russian. Vestnik Leningradskogo Universiteta 3 (15): 38–52.

- Sokolovskaya A. P. 1972. Karyological characteristics of the flora of the Leningrad district (in Russian). *Vestnik Leningradskoza Zbuversutetam* 21: 60.
- Solereder H. & Meyer F. J. 1933. Systematische Anatomie der Monokotyledonen, I, pt. 1: 50–67. Gebrüder Borntraeger, Berlin.
- Stace C. A. 1975. Hybridization and the flora of the British Isles. Academic Press, London, New York, San Francisco. pp. 1–626.
- Taylor R. L. & Mulligan G. A. 1968. Flora of the Queen Charlotte Islands. Part 2. Cytological aspects of the vascular plants. Queen's Printer, Ottawa. pp. 148.
- Tutin T. G., Heywood V. H., Burges N. A., Moore D. M., Valentine D. H., Walters S. M. & Webb D. A. 1980. Flora Europeae 5: 274–275. 1980. Cambridge Univ. Press, Cambridge.
- Váčová M. in Májovský J. 1976. Index of chromosome numbers of Slovakian flora (Part 5). *Acta Fac. Rerum Nat. Univ. Comenianae, Bot.*, 25: 1–18.
- Velenovský J. & Viníklář L. 1926. Flora cretacea Bohemiae. I. *Rozpr. Stat. Geol. Ust. ČSR.* 1: 1–157.
- Waggers Nielsen L., Nielsen K. & Sand-Jensen K. 1985. High rates of production and mortality of submerged *Sparganium emersum* Rehman [sic] during its short growth season in a [sic] eutrophic Danish stream. *Aquatic Botany* 22: 325–334.
- Wulff H. D. 1938. Chromosomenstudien an der schleswig-holsteinischen Angiospermen-Flora. II. *Ber. Deutsch. Bot. Ges.* 56: 247–254.
- Yuzepchuk S. V. 1934. In Komarov V. L., Flora of the USSR. Izdatel'stvo Akademii Nauk SSSR. (Leningrad) 1: 216–229.
- Zhukova P. G. 1980. Chromosome numbers of some Southern Chukotka plant species (in Russian). *Botaničeskij Žurnal Akad. Nauk SSSR. (Leningrad)* 65 (1): 51–59.
- Zhukova P. G. & Petrovsky V. V. 1976. Chromosome numbers of some Western Chukotka plant species, II (in Russian). *Botaničeskij Žurnal Akad. Nauk SSSR. (Leningrad)* 61 (7): 963–969.
- Zhukova P. G. & Tikhonova A. D. (1971). Chromosome numbers of certain plant species indigenous to the Chukotskiy province (in Russian). *Botaničeskij Žurnal Akad. Nauk SSSR. (Leningrad)* 56: 868–875.


Maps




Map 1. *Sparganium hyperboreum* in Eurasia. Open circles denote unconfirmed literature records.




Map 2. *Sparganium hyperboreum* in N. America, including records published by Harms (1973).




Map 3. *Sparganium natans* in Eurasia. Open circles denote hybrid *S. hyperboreum × S. natans*.




Map 4. *Sparganium natans* in N. America, including records published by Harms (1973). Open circles denote hybrid *S. hyperboreum* × *S. natans*.




Map 5. *Sparganium angustifolium* in Eurasia. Open circle denotes unconfirmed literature record.




Map 6. *Sparganium angustifolium* in N. America.



Map 7. *Sparganium glomeratum* in Eurasia.




Map 8. *Sparganium gramineum*.




Map 9. *Sparganium angustifolium* × *S. gramineum*.


Map 10. *Sparganium emersum* subsp. *emersum* × *S. gramineum*.



Map 11. *Sparganium fluctuans*, closed circles; *S. glomeratum*, closed triangles confirmed record, open triangles unconfirmed literature records.



Map 12. *Sparganium emersum* subsp. *emersum* in Eurasia.



Map 13. *Sparganium emersum* in N. America: subsp. *emersum* closed circles; subsp. *acaule* open circles.

A monographic study of the genus *Sparganium* (Sparganiaceae). Part 2, Subgenus *Sparganium*

Autor(en): **Cook, Christopher D.K. / Nicholls, Marc S.**

Objekttyp: **Article**

Zeitschrift: **Botanica Helvetica**

Band(Jahr): **97(1987)**

Heft 1

Erstellt am: **Apr 28, 2013**

Persistenter Link: <http://dx.doi.org/10.5169/seals-67851>

Nutzungsbedingungen

Mit dem Zugriff auf den vorliegenden Inhalt gelten die Nutzungsbedingungen als akzeptiert. Die angebotenen Dokumente stehen für nicht-kommerzielle Zwecke in Lehre, Forschung und für die private Nutzung frei zur Verfügung. Einzelne Dateien oder Ausdrucke aus diesem Angebot können zusammen mit diesen Nutzungsbedingungen und unter deren Einhaltung weitergegeben werden. Die Speicherung von Teilen des elektronischen Angebots auf anderen Servern ist nur mit vorheriger schriftlicher Genehmigung des Konsortiums der Schweizer Hochschulbibliotheken möglich. Die Rechte für diese und andere Nutzungsarten der Inhalte liegen beim Herausgeber bzw. beim Verlag.

SEALS

Ein Dienst des *Konsortiums der Schweizer Hochschulbibliotheken*
c/o ETH-Bibliothek, Rämistrasse 101, 8092 Zürich, Schweiz
retro@seals.ch
<http://retro.seals.ch>

A monographic study of the genus *Sparganium* (Sparganiaceae).

Part 2. Subgenus *Sparganium*

Christopher D. K. Cook and Marc S. Nicholls

Institut für Systematische Botanik der Universität Zürich, Zollikerstr. 107, CH-8008 Zürich,
Switzerland

Manuscript accepted January 9, 1987

Abstract

Cook, C. D. K. & Nicholls, M. S., 1987. A monographic study of the genus *Sparganium* (Sparganiaceae). Part 2. Subgenus *Sparganium*. *Bot. Helv.* 97: 1–44.

A taxonomic revision of *Sparganium* (Sparganiaceae) subgenus *Sparganium* is presented. Each species has a full description, short diagnosis, list of synonyms with typifications, distribution map with notes on ecology, variation and hybrids. Seven species and five subspecies are recognized. Two new names are used: *Sparganium erectum* subsp. *stoloniferum* (F. Hamilton ex Graebner) C. D. K. Cook & M. S. Nicholls *comb. nova*; *S. eurycarpum* subsp. *coreanum* (Léveillé) C. D. K. Cook & M. S. Nicholls *comb. nova*. An index to the complete genus *Sparganium* is given.

Introduction

Part 1 of this study is published in *Botanica Helvetica*, 96: 213–267 (1986) and includes introductory chapters for the entire genus: materials and methods, special terminology, family and generic description with typification, distribution, fossils, anatomy and morphology of vegetative and floral parts, chromosome numbers, floral biology, dispersal, ecology, parasites and animal feeders, applied aspects and a key to all species and subspecies. Individual accounts of the species and subspecies of subgenus *Xanthosparganium* are also included in Part 1.

This part of the study gives individual accounts of the seven species and five subspecies comprising subgenus *Sparganium*. An index to the genus is also provided on p. 30.

Acknowledgement

We would like to thank the Fonds National Suisse de la Recherche Scientifique (Nr. 3.390-0.83) for a post-doctoral fellowship for Marc S. Nicholls. In addition to the people acknowledged in part 1 (*Bot. Helv.* 96: 214) we would also like to thank Dr. N. Andreev, Sofia, Bulgaria and Dr. J. M. Montserrat, Barcelona, Spain.

Species of subgenus Sparganium

Perianth segments not translucent with a thickened dark-brown to black pad of tissue near the apex, apex emarginate to entire or nearly so.

Key to species in *Botanica Helvetica* 96: 228–230, 1986.

8. *Sparganium fallax* Graebner ex Graebner in Engler, *Pflanzenreich* 2 (IV. 10): 15. September 1900 – *S. fallax* Graebner, *Allg. Bot. Zeitschr.* 4: 32. February 1898, nomen sub-nudum. Type: “Asien: von Sikkim! Khasia! und Ostbengalen bis Japan” (holotype: Bangla Desh, “East Bengal, Herb. *Griffith* 5618, specimen with label in Graebner’s handwriting dated 5 Jan. 1898, B; numerous syn- or isotypes in E, K)
- = *S. yamatense* Matsumura ex Makino in Hara, *J. Jap. Botany* 12 (3): 171. March 1936 – *S. yamatense* Matsumura, *Indig. Pl. Jap.* (2) 1: 24. 1905, nomen nudum. Type: Japan, prov. Yamato, Nara, 15 July 1883, *J. Matsumura* (holotype: TI)
 - = *S. confertum* Y.-D. Chen, *Acta Phytotaxonomica Sinica* 19 (1): 53. 1981. Type: China, Yunnan, Gaoligong Shan, alt. 3100 m, 11 October 1940, *K. M. Feng* 8379 (holotype: Acad. Sin., probably PE n.v.; paratype: 17 September 1938, *T. T. Yü* 20343, E).
 - = *S. yunnanense* Y.-D. Chen, *Acta Phytotaxonomica Sinica* 19 (1): 54. 1981. Type: China, Yunnan, Menghai Xian, alt. 1530 m, May 1936, 玉肩毛 [Z. W. Wang] 73555 (holotype: Acad. Sin., probably PE n.v.).

Robust to slender plants with erect, emergent or occasionally floating leaves. Stolons up to 25 cm or more long, ca. 2–3.5 mm diameter.




Fig. 13. Diagrammatic representation of the inflorescences of: a, *Sparganium fallax*; b, *S. subglobosum*; c, *S. japonicum*.

Basal leaves usually erect and partially emergent, 40–60(–80) cm long, usually exceeding the stem, (4–)5–10(–15) mm wide, sheathing but not inflated at base, carinate (usually to apex); apex rounded and dark-tipped.

Flowering stems 30–60 cm long, usually erect, simple, with 2–4, elongated, sterile nodes below the inflorescence.

Inflorescence bracts erect, diverging from the main axis at an acute angle, not or slightly inflated at base, sometimes with fine hyaline margins; lowermost bract up to ca. 20–35(–60) cm long, 1–2 or more times as long as the inflorescence (Fig. 13 a).

Female heads 3–4(–6), usually supra-axillary, sometimes extending beyond the next internode (Fig. 14 a), usually remote with zig-zag main axis, sessile above and usually pedunculate below; in fruit (10–)15–20(–34) mm diameter; pendule up to ca. 3 cm long.

Male heads 5–8 or more, 7–18 mm diameter, separated from the uppermost female head by a (8–)10–15(–50) mm long internode; the lower ones usually remote becoming overlapped above; the lowest sometimes supra-axillary and subtended by a leaf-like bract up to 6 cm long but usually less (the male inflorescence thus may appear to be stalked).

Female flowers with perianth segments oblong to spatulate, distinctly united below and sometimes partially united above (Fig. 14 a) and then irregularly shaped, ca.




Fig. 14. Perianth segments, fruits and endocarps of: a, *Sparganium fallax*; b, *S. subglobosum* (scale bar: 1 mm).

3.5–5.5 mm long, more than half as long as the fruit, inserted at top half of pedicel, apex and midrib dark; pedicel ca. 1–3 mm long; stigmas (0.4–)0.8–1.2(–1.6) mm long.

Male flowers with filaments up to 6 mm or more long; anthers (0.8–)1.0–2.0(–2.2) mm long.

Fruits fusiform, 5–6 mm long, sometimes constricted around the middle, ca. 2–2.5 mm wide, light brown, dull, tapering above to a 1.6–2.5(–3.0) mm long beak, tapering below into a 1.5–3 mm long pedicel; endocarp ovate ca. 3 mm long and 2 mm wide, truncate at micropyle, tapered below into an up to ca 3 mm long stalk (Fig. 14a).

Diagnostic features

Inflorescence simple; perianth segments with dark tips and midribs; female heads remote, mostly supra-axillary, some sometimes concaulescent beyond the next internode (Fig. 13a); leaves carinate to apex, usually more than 5 mm wide; lower bract 1–2 times as long as inflorescence; fruit beak more than 1.6 mm long; stigmas less than 1.6 mm long (see Table 4). It sometimes superficially resembles *S. emersum* but can be distinguished by the perianth segments with dark tips and shorter stigmas.

Distribution

S. fallax is an eastern Asian species extending from central Honshu, Japan, south to Kyushu, through southern China westwards through Yunnan and northern Burma to south eastern Tibet, Assam and Meghalaya, India, Map 14. It is also found in the highlands of New Guinea and reported in the mountains of central Sumatera (usually misidentified as *S. simplex*).

Ecology

We have no first-hand information. From herbarium specimens it appears to be an emergent aquatic growing in shallow water. In the north of its range it grows at sea level in the south at about 2000 m in New Guinea.

Variation

The variation observed on herbarium specimens is incorporated in the formal taxonomic description.

Hybrid

8×9. *S. fallax*×*S. subglobosum*

This hybrid has been reported but we doubt that it exists, see p. 8.

9. *Sparganium subglobosum* Morong, Bull. Torrey. Bot. Club 15 (3): 81. Pl. 79, Fig. 1. 2 March 1888. Type: New Zealand, Bay of Islands, U.S. Exploring Expedition under Command of Capt. Wilkes (holotype; GH n.v.)

= *S. stenophyllum* Maximovicz ex Meinshausen, Bull. Soc. Imp. Nat. Moscou N.S. 3(1889): 171. 1890. Type: USSR., Primorskiy Kray, Ussuriysk (Voroshilov), "S.O. Mandschurien, in einem kleinen Landsee beim Dorf Nikolsk (Goldenstädt), 28. Juli 1872", Maximovicz (holotype: LE not found).

= *S. antipodum* Graebner, Allg. Bot. Zeitschr. 4 (2): 33. Febr. 1899. Type: New Zealand or Australia, „Neu-Seeland: verbreitet. – Australien: Victoria, F. v. Mueller” “Fruc-

Table 4. Comparison of *Sparganium emersum* and the species of the *S. americanum* group

	<i>emersum</i>	<i>fallax</i>	<i>subglossum</i>	<i>japonicum</i>	<i>americanum</i>	<i>androcladum</i>
Perianth segment apex	translucent	dark	dark	dark	dark	dark
Leaves: width (mm)	(2-)4-12(-18)	(4-)5-10(-15) keeled	2-4(-9) keeled	(3-)5-10(-20) triangular-flat	(5-)7-12(-18) keeled-triangular flat	5-12 keeled triangular-flat
Leaves: T.S. proximal	triangular (- flat)	keeled-triangular	keeled-triangular	flat	flat	triangular-flat
Leaves: T.S. distal	erect (floating)	erect	erect	ascending	spreading	ascending
Lowest bract: position	1-4× longer	1-2× longer	shorter-equal	1-2× longer	1-2× longer	longer
Lowest bract: relation to inflorescence	no branch	no branch	♀ + ♂ or ♂ only	no branch	♀ + ♂ or simple	♂ only or simple
Inflorescence: lowest branch	remote	very remote	remote	upper approximate	remote	remote
Female heads: remote or approximate	(approximate)	always	never	very rare	very rare	rare
Female heads: supra-axillary	usually					
Fruiting head: diameter (mm)	16-25(-35)	(10-)15-20(-34)	12-16(-23)	15-20	15-25	25-35
Fruit: texture	shiny	dull	shiny	dull	dull & pitted	shiny above dull pitted below
Fruit: body length (mm)	3.5-5.5	5-6	4-5(-6.5)	(-3)4-6	(3.5)-4-5(-5.5)	5-7
Fruit: body diameter (mm)	1.8-2.5	2-2.5	2-3	ca. 2	ca. 2	2.5-3
Fruit: beak length (mm)	2-4.5(-6)	1.6-2.5(-3)	1-2	1.5-3	(1.5)-3-4(-5)	4.5-7
Fruit: pedicel length (mm)	1-4	1.5-3	0-1	2-5	1-2(-3)	2.4-4
Stigma length (mm)	(1-)1.5-2(-2.5)	(0.4)0.8-1.2 (-1.6)	0.6-1.0(-1.2)	1-1.5	(0.9-)1-2(-3)	2-4
Endocarp length (mm)	2.5-3.5	ca. 3	2.5-3	ca. 3.5	3.5-4	ca. 4.5
Anther length (mm)	1-1.5(-2)	(0.8-)1-2(2.2)	0.5-0.9(-1)	0.7-1.0	0.8-1.3	1-1.6

- tus mihi ignoti" (lectotype: chosen here, New Zealand, "Nov. Zeland", s.d., *J. D. Hooker s.n.* det. Graebner 5 Jan. 1898 in B; isolectotype K; paratype K).
- = *S. nipponicum* Makino ex Makino. *J. Jap. Bot.* 3 (6): 21. 1926. — *S. nipponicum* Makino, *Bot. Nat. Fr. Auth. Priv. Cabinet Fl. Jap.* 2: 40. 1898, nomen seminudum. Type: Japan, Honshu, "prov. Shimousa, Junsai-numa, prope Ichikawa, September 1893." T. Makino (holotype: MAK n.v., there is material in MAK collected at this locality by Makino in 1894 but none has been found collected in September 1893; material in TI collected by Makino from this locality in September 1893 is the holotype of *S. macrocarpum*, see p. 28).
- = *S. limosum* Y.-D. Chen, *Acta Phytotax. Sinica* 19 (1): 52. February 1981. Type: China, Yunnan, Goaligong Shan, alt. 1750 m, 26 June 1938, T. T. Yu 19197 (holotype: Acad. Sin. probably PE n.v.)

Robust to slender plants with erect and emergent or rarely floating leaves. Stolons slender up to 20 cm or more long, 2–3 mm diameter.

Basal leaves usually erect and partially emergent or rarely floating, (25–)30–60(–120) cm long, usually exceeding the inflorescence, (1–)2–4(–9) mm wide, sheathing but not inflated at base, carinate (usually to apex), glossy, with a rounded blackish apex.

Flowering stems (16–)30–50(–130) cm long, usually erect, emergent, simple or with 1 or 2 branches at base of inflorescence, lateral branches axillary with (0–)1–2 female heads below and 3–4 male heads above (in Australasia some specimens lack female heads on branches); main inflorescence axis with (1–)2–3(–4) female and (3–)6–9(–11) male heads (Fig. 13 b).

Inflorescence bracts more or less erect, not or very slightly inflated at base, sometimes with very fine hyaline margins; lowermost bract up to ca. 15 cm long, about half as long as or equalling but very rarely exceeding the inflorescence.

Female heads axillary, sessile or very rarely with a peduncle less than 5 mm long, remote (each distinct and not touching), in fruit 12–16(–23) mm diameter.

Male heads 9–13 mm diameter, separated from the uppermost female head by a (5–)10–16(–22) mm long internode, usually distinct but not remote from each other at anthesis, usually ebracteate.

Female flowers with perianth segments oblong to spatulate, with a dark brown tip and midrib, not united below, at maturity one-third to half as long as the fruit; sessile or with an up to 1 mm long pedicel; stigmas 0.6–1.0(–1.2) mm long, lanceolate (Fig. 14 b).

Male flowers with filaments 2.5–3.2(–3.8) mm long; anthers 0.5–0.9(–1.0) mm long.

Fruits ovoid to almost globose, 4–5(–6.5) mm long, ca 2–3 mm wide, yellowish to pale brown, shiny-surfaced, tapering and angular in transverse section below, sessile or with up to 1 mm long pedicel, narrowed above to conic-based summit with a blackish, 1–2 mm long beak; endocarp ovoid to pyriform, 2.5–3 mm long and 1.6–1.8 mm wide, almost smooth with very faint longitudinal veins, stalk not persistent (Fig. 14 b).

Diagnostic features

Perianth segments with brown tips and midribs; leaves rarely more than 5 mm wide and keeled to the apex; lowest inflorescence bract not or rarely exceeding the inflorescence; inflorescence usually with 1 or 2 branches; all female heads axillary and remote; fruits ovoid to almost globose, shiny-surfaced, yellowish to pale-brown a with blackish beak; stigmas rarely exceeding 1 mm long (see Table 4).

Distribution

S. subglobosum is an eastern species with a somewhat disjunct distribution (Map 15). In the northern hemisphere it is found in Manchuria, north-west China and Japan extending from Hokkaido southwards to Okinawa. It reappears in south-west China in Yunnan (*Maire* 896 in E, and see *Chen Yao-Dong*, 1981) and in Vietnam. There is a single gathering from Megalaya, India ("Khasia, c. 2000 m, 26 August 1850, *Hooker fil.* & *Thompson*, E, K and other herbaria). In the southern Hemisphere it occurs in New Zealand, Australia and the highlands of new Guinea. In New Zealand it is found in the North Island extending southwards into the South Island at about 42°S with isolated records from Canterbury. In Australia it is found along the eastern coast from S.E. Queensland to Victoria near Melbourne, in coastal to highland regions from sea-level to 1300 m altitude.

Ecology

S. subglobosum is a plant found in still or slowly-flowing water less than 1 m deep on the margins of creeks, rivers and lakes. In Australia it is characteristically in ditches, swamps and creek edges in dry sclerophyll forest and cleared grazing land. It is rather indifferent to soil, being found on sand, loam, peat and red soils overlying basalt, sandstone, granite and quartz porphyry. The following species are commonly associated with *S. subglobosum*: *Eleocharis sphacelata* R. Br., *Glyceria australis* C. E. Hubbard, *Myriophyllum propinquum* A. Cunn. s.l., *Philydrum lanuginosum* Blanks & Sol. In Australia it is often eaten by stock during the dry season and it is claimed to prevent the erosion of banks.

In Australia it flowers mostly from December to February and sets fruit by May or June. In Japan it flowers mostly from June to August and the fruit ripens by October.




Fig. 15. *Sparganium subglobosum*, comparison of plants from New Zealand and Australia (▲) with plants from Japan (●). Vertical axis: ratio of the length of the lowest bract to the length of the inflorescence; horizontal axis: leaf-width.

Variation

S. subglobosum plants are typically slender and erect with leaves rarely more than 5 mm wide. Occasionally more robust specimens are encountered with wider leaves (reaching a maximum of 9 mm) but they do not differ in other respects. One specimen (Australia: NSW., Port Macquarie, November 1897, J. H. Maiden 2404 – NSW) has floating leaves up to about 120 cm long; this specimen was seen by Rothert who suggested the plant had been growing in exceptionally deep water.

S. subglobosum has been little studied previously and there are areas of its range, particularly India, south-west China and New Guinea, where it remains poorly collected. Nevertheless, the degree of intraspecific variation observed in our study is far less than that of most other *Sparganium* species.

A comparison of plants from Japan with those from Australia and New Zealand revealed minor differences in leaf width and bract/inflorescence length ratio (see Fig. 15). Japanese plants have slightly narrower leaves (rarely exceeding 4 mm) and the lowermost bract tends to be shorter than the inflorescence while Australian plants mostly have wider leaves and slightly longer bracts. There is, however, considerable overlap in both characters and we have seen very little material outside Japan and Australia. We see no grounds for formally recognizing any intraspecific taxonomic categories.

Hybrid

8×9, *S. fallax* × *S. subglobosum*

On an herbarium sheet in Kew bearing specimens of both species from Khasia (Meghalaya, India), Rothert suggests one of the specimens is of hybrid origin. We contend that the specimen is in fact a slender plant of *S. fallax* and doubt that this hybrid exists.

10. *Sparganium japonicum* W. Rothert in Fedtschenko, Flora Aziatskoj Rossii 1: 26, Tab. 2. 1913. Type: USSR, Primorskiy Kray, "southern Ussuri Region, Village Grigor'evka, on the River Čichc, 1888, Pal'čevskij" (holotype: LE n.v.).

Slender to robust plants usually erect and emergent or floating in deeper water. Stolons up to ca. 15 cm long, 2.5–3.0 mm diameter.

Basal leaves erect or floating, up to 80 cm or more long usually exceeding the inflorescence, (3–)5–10(–20) cm wide, carinate or flattened below, usually flattened above, without keel; apex rounded and dark; midrib obscure towards apex.

Flowering stems ca. 40–80 cm long; inflorescence up to ca. 20 cm long, simple.

Inflorescence bracts spreading or spreading-ascending, not inflated at base, usually flattened and not keeled; lowermost bract ca. 12–40 cm long, 1–2 times as long as the inflorescence; upper bracts spreading to reflexed (Fig. 13 c).

Female heads (2–)3–6, axillary, the lower 1–3 often long-pedunculate, the upper heads usually approximate in fruit; in fruit ca. 15–20 mm diameter.

Male heads 5–9, the lower ones usually remote the upper ones usually approximate, ca. 10 mm diameter, separated from the uppermost female head by an internode up to 15 mm long.

Female flowers with perianth segments oblong to lanceolate or spatulate, with a dark brown tip and midrib, 4–5 mm long, more than half as long as fruit, attached to top half of pedicel; pedicel 2–5 mm long; stigmas ca. 1–1.5 mm long.

Male flowers with dark tipped perianth segments; filaments ca. 4–5 mm long; anthers ca. 0.7–1.0 mm long.

Fruits fusiform (3–)4–6 mm long, ca. 2 mm wide, sometimes constricted around the middle, no obvious difference between upper and lower halves, brown to dark brown, dull-surfaced, tapering below into an obconic base with a 2–5 mm long pedicel; above gradually narrowed into a straight beak, ca. 1.5–3 mm long; endocarp ovoid, ca. 3.5 mm long, ca. 2.5 mm diameter, with an up to 3–5 mm long stalk, smooth or with very fine longitudinal veins (Fig. 16 a).

Diagnostic features

Perianth segments with dark brown tips and midribs. Like *S. americanum* (see p. 10) but inflorescence always simple; upper female heads approximate; fruit lacking dark brown glands, not pitted; beak not exceeding 3 mm; pedicel usually 3 mm or more long. Like *S. fallax* (p. 2) but usually all heads axillary; leaves flat at apex; upper bracts spreading to reflexed; upper female heads approximate; main axis not zig-zag; fruit dark-brown. Like *S. subglobosum* (p. 4) but fruits fusiform; inflorescence simple, fruits dark brown, dull; leaves usually more than 5 mm wide and distally flat; stigmas exceeding 1 mm long; pedicel usually exceeding 1 mm persistent in fruit; endocarp exceeding 3 mm long, stalked at maturity (see Table 4).




Fig. 16. Perianth segments, fruits and endocarps of: a, *Sparganium japonicum*; b, *S. americanum*; c, *S. androcladum* (scale bar: 1 mm).

Distribution

S. japonicum is from E. Asia and shows a compact distribution pattern. It is found in Japan from northern Honshu to southern Kyushu and in southern Primorskiy Kray (north of Vladivostok) and in Korea, see Map 16.

Ecology

We have no first-hand information. From herbarium material it appears to be a perennial, emergent aquatic growing in shallow water. On one herbarium sheet (Okinawa Island, 18 May 1955, *Hatusima* 17886, TI) it is reported to grow as an annual in a wet field – we consider this rather unlikely and suggest that the specimen might be incorrectly labelled. We have not included this locality on Map 16.

Variation

S. japonicum shows close affinities to *S. americanum* and could well be recognized at an infraspecific rank. Without more detailed work including field studies we prefer to keep them as separate species rather than to make a new nomenclatural combination. Its identity is clear.

11. ***Sparganium americanum*** Nuttall, Genera North American Plants 2: 203. 14 July 1818 = *S. simplex* var. *americanum* (Nuttall) Engler in Engler & Prantl, Pflanzenfamilien 2 (1): 192. 1887. Type: USA, Pennsylvania, "vicinity of Philadelphia", Nuttall (holotype: PH n.v., ? BM).
- = *S. americanum* var. *rigidum* Clausen, Rhodora 39: 189. May 1937. Type: USA, New Jersey, Ocean Co., Tuckerton Creek Pond, 22 September 1934, Edwards & Clausen (holotype: GH).
 - *S. simplex* var. *nuttallii* Engelmann, in A. Gray. Manual Botany Northern U.S., ed. 5, 481. 1867 nomen illeg., based on *S. americanum* Nuttall.

Slender to robust plants, erect and emergent or floating in deeper water. Stolons 10–20 cm or more long, 3–4 mm diameter.

Basal leaves erect or floating, rarely exceeding the inflorescence, (30–)40–100 cm long, (5–)7–12 mm wide, flattened or carinate below, flat above; apex rounded and dark; midrib obscure.

Flowering stems up to ca. 100 cm long; inflorescence 10–25(–30) cm long, with up to 3 branches or occasionally simple (Fig. 17 a); branches axillary with 1–3 female and up to 7 male heads; main axis with 2–4 female and 4–10 male heads (Fig. 17 b).

Inflorescence bracts spreading or spreading-ascending, sometimes inflated at base, usually not keeled; lowermost bract (5–)10–35(–40) cm long, 1–2 times as long as the inflorescence.

Female heads axillary, usually sessile, all remote, in fruit ca. 15–25 mm diameter.

Male heads more or less distinct but not all remote at anthesis, 10–15(–18) mm diameter, separated from the uppermost female head by a (4–)10–20(–26) mm long internode.

Female flowers with perianth segments oblong to oblanceolate or spatulate, with dark brown tip and midrib often dotted with brown glands (Fig. 16 b), about half to three-quarters as long as the fruit, attached at top of pedicel; pedicel up to 3 mm long; stigmas (0.9–)1–2(–3) mm long.




Fig. 17. Diagrammatic representation of inflorescences of: a, b, *Sparganium americanum* (a, unbranched; b, branched); c, *S. androcladum*.

Male flowers with dark-tipped perianth segments; filaments 4–5 mm long; anthers 0.8–1.3 mm long.

Fruits fusiform, (3.5–)4–5(–5.5) mm long, ca. 2 mm wide, sometimes slightly constricted around the middle, brown to dark brown, dull-surfaced and sometimes pitted, tapering below into an obconic base with a 1–2(–3) mm long pedicel; above rather abruptly narrowed into a straight or curved but not hooked beak, (1.5–)3–4(–5) mm long; endocarp ovoid, 3.5–4 mm long, ca. 2 mm diameter, smooth with fine longitudinal veins (Fig. 16 b), very difficult to remove from fruit.

Diagnostic features

Perianth segments with dark tips and midribs; inflorescence simple or lowest branch bearing both female and male heads; female heads axillary, in fruit not exceeding 25 mm diameter; fruits brown to dark-brown with surface dull and pitted, rarely more than 5 mm long and 2 mm diameter with a straight beak not more than 5 mm long; stigmas rarely more than 2 mm long; leaves usually more than 5 mm wide, usually keeled or carinate below but flattened above; inflorescence bracts spreading, the lowest longer than the inflorescence; pedicel of female flowers or fruits rarely more than 2 mm long (see Table 4). *S. americanum* shows close affinities to *S. japonicum* (p. 8) and *S. androcladum* (p. 12).

Distribution

S. americanum is widespread in North America ranging from Newfoundland west to Ontario and North Dakota, extending south to northern Florida, southern Alabama,

Louisiana, Oklahoma and Texas, with an isolated record just north of 23°N. In the highlands of Mexico (Durango Prov., El Salto, 2600 m, 7 July 1956, *Dickerman 1005*, GH), see Map 17.

Ecology

S. americanum is found in shallow, still or flowing water in a variety of natural and artificial aquatic habitats. It is usually found near the shores and may build largish stands. In the Northeast it is a characteristic species of the bank vegetation along silted streams and is found in marshes.

Variation

S. americanum is a polymorphic species which has been studied in the southeastern USA by Beal (1960). Of particular interest to Beal was the correlation and distribution of the following characters: stem height, leaf width, thickness and texture, degree of branching of the inflorescence, position of the female heads, size of the fruit and length of style and stigma. On the basis of these characters Beal (1960) recognized three races: (1) the Coastal Race, with stigmas 1.5 mm or more long, relatively wide leaves and 2–5 inflorescence branches (see Fig. 17 b); (2) the Appalachian Race, with stigmas 0.9 mm or shorter, narrower leaves and simple or sparingly branched inflorescences (Fig. 17 a), and (3) the Ubiquitous Race, based on plants intermediate between the first two races. As the names suggest, the first two races are geographically somewhat restricted while the third race is found throughout the range. Between all three races there is considerable overlapping and therefore Beal (1960) found it "inappropriate to dignify these races with formal names."

Beal (1960) made detailed morphological studies of 148 collections from the Carolinas. He attempted various correlations and designed an index. Despite all this playing around with numbers, the single and most frustrating fact to emerge about *S. americanum* is that, while undoubtedly polymorphic, it does not fit well into any of those patterns favoured by students of biosystematics and the like. Beal explored three explanations of the distribution patterns observed: environmental effects, introgressive hybridization and incipient race-formation. Having demolished the second of these he came to the sensible (if banal) conclusion that "a complex of interacting genetical and ecological factors have been involved." His subsequent call for further experimental work involving transplant and hybridization studies remains to be taken up.

Variation in *S. americanum* has produced an attendant nomenclatural confusion. In the original description of *S. americanum*, Nuttall (1818) referred only to plants of what Beal (1960) called the Appalachian Race. Unfortunately Morong (1888) followed by Mohr (1910), Fernald & Eames (1907) and Rydberg (1909) misinterpreted robust plants (Beal's Coastal Race) and failed to distinguish them from *S. androcladum* (see below). Fernald (1922) cleared up these problems, reinstating the species *S. androcladum*.

Clausen (1937) recognized a new variety *rigidum* for plants having narrow, stiff leaves and a supra-axillary lowest female head. Beal's (1960) and our own studies suggest these features are not correlated, are scattered throughout the range of *S. americanum* and are not worthy of taxonomic recognition.

12. ***Sparganium androcladum* (Engelmann)** Morong, Bull. Torrey Bot. Club 15 (3): 78.
2 March 1888 = *S. simplex* var. *androcladum* Engelmann in A. Gray, Manual

- Botany Northern U.S., ed. 5, 481. 1867 = *S. americanum* Nuttall var. *androcladum* (Engelmann) Fernald & Eames, Rhodora 9: 87. May 1907. Type: USA, "New England" Engelmann (holotype: MO). Unfortunately Morong (op.cit.) misinterpreted robust plants of *S. americanum* and failed to distinguish them from Engelmann's var. *androcladum* (op.cit.). Morong's error was perpetuated by Mohr (Contr. U.S. Nat. Herb. 6:327. 1901) and Rydberg (N. Amer. Flora 17: 8. 1909). Fernald & Eames (op.cit.) proposed the new combination (*S. americanum* var. *androcladum*) and applied it to robust pants of *S. americanum*; the species now known to be *S. androcladum* they described as a new species, *S. lucidum*. Fernald (Rhodora 24: 26. February 1922) cleared up these problems, reinstating *S. androcladum* in the correct sense and reducing *S. lucidum* to synonymy.
- = *S. lucidum* Fernald & Eames, Rhodora 9: 87. May 1907. Type: USA, Massachusetts, "small pond Medford, 29 July 1860, Wm. Boott (lectotype: GH).
 - *S. simplex* Hudson var. *androgyna* Meinshausen, Bull. Acad. Imp. Sci. St. Pétersbourg N.S. 4 (36): 31. December 1893. This is probably a misprint for "*androcladum*".

Robust erect or rarely floating plants. Stolons up to 20 cm or more long, up to 4 mm diameter.

Basal leaves usually erect and emergent, up to ca. 100 cm or more long, 5–12 mm wide, exceeding the inflorescence, distinctly keeled from base to at least the middle; apex rounded and dark; midrib prominent and thickened at apex.

Flowering stems up to 100 cm or more long, rigid and erect; inflorescence normally branched or occasionally simple; main axis bearing (1–)2–4, sessile or at base shortly pedunculate, axillary female heads and up to 8 or more male heads; lateral branches axillary, the lowest with 1–6 male heads and nearly always without female heads or very rarely with 1 female head, the second or third lowest rarely bearing 1 female head otherwise with 1–6 male heads (Fig. 17 c).

Inflorescence bracts ascending, not inflated below, distinctly triangular in transverse section, often keeled; the lowest ca. 25–40 cm long, exceeding the inflorescence; bracts subtending upper female and lower male heads usually green and more or less leaf-like.

Female heads axillary or rarely supra-axillary, sessile or sometimes the lowest shortly pedunculate, all remote, usually confined to main axis, rarely the second or third branch bearing solitary heads below the males, in fruit 25–35 mm diameter.

Male heads ca. 10–15 mm diameter, distinctly separated from female ones by a (4–)10–20(–26) cm long internode, the lower ones remote and distinct from each other.

Female flowers with perianth segments oblong to oblanceolate or spatulate, with dark brown tip and midrib, attached to top of pedicel, half to two-thirds as long as the body of the fruit; pedicel 2.4–4 mm long; stigmas 2–4 mm long.

Male flowers with dark-tipped perianth segments; filaments up to ca. 6 mm long; anthers ca. 1–1.6 mm long.

Fruits ovoid to fusiform, 5–7 mm long, 2.5–3 mm diameter, often constricted around the middle; the lower part brown, almost terete and dotted with dark reddish-brown glands, with an obconic base and persistent pedicel; the upper part smooth, shiny, brown to olive-brown, gradually tapered above into 4.5–7 mm long beak often hooked at apex; pedicel 2.4–4 mm long, endocarp ovoid, ca. 4.5 mm long, ca. 2.5 mm diameter, smooth with the fine longitudinal veins (Fig. 16 c), very difficult to remove from fruit.

Diagnostic features

Like *S. americanum* (p. 10) but inflorescence usually branched, the lowest or lower branches usually bearing male heads only or very rarely single female heads; fruiting heads more than 25 mm diameter, usually sessile and axillary or rarely one supra-axillary; leaves and bracts usually carinate to apex, usually more than 5 mm wide; fruits shiny above and dull, pitted and glandular below, 5–7 mm long, 2.5–3 mm diameter, with usually curved beak 4.5–7 mm long; pedicel of female flowers usually exceeding 2 mm long; endocarp usually more than 4 mm long; stigma 2–4 mm long; anthers 1–1.6 mm long.

Distribution

The distribution of *S. androcladum* has been studied in detail by Beal (1960). It is a species confined to northeastern North America. Its range extends from southern Ontario and Quebec southwards to southern Virginia, then westwards through northern Tennessee to Missouri (Map 18). Rather strangely, it seems to be absent in the middle of the range and is not known from Michigan, Indiana and Kentucky and is rare in Ohio and West Virginia.

Ecology

We have no first-hand knowledge of the ecology of *S. androcladum*. It appears to be a plant of shallow water growing in marshes or along the shores of lakes, rivers and streams.

Variation

S. androcladum is a remarkably constant species. It shows close affinities with *S. americanum* and it is sometimes difficult to distinguish the two when the plants are immature (see Table 4).

13. *Sparganium erectum* L. sensu lato

- = *S. ramosum* forma *simplicior* Rothert, Acta Horti Botanici Univ. Imper. Jurevensis 11: 19, 1910. *Type*: no specimen cited; used for unbranched plants of *S. erectum*.
- *S. barrerae* Sennen, Bull. Soc. Bot. France 74: 404, 1927 nomen nudum

Robust erect, emergent or rarely floating or submerged herbs. Stolons up to 60 cm long, (3–)6–9 mm diameter. Corms 1.5–4.5 cm diameter, when mature hard and woody, each bearing up to 9 stolons.

Basal leaves obliquely erect and fan-like, usually emergent, often pink at base, (30–)50–150(–350) cm long, (6–)10–20(–28) mm wide, carinate from base to apex, with 3–8 layers of gas chambers.

Plants in very deep or swiftly flowing water may develop submerged or partly floating leaves but usually remain sterile.

Flowering stems erect, 20–100(–150) cm long, nearly always branched, with 1–2(–4) sterile nodes between the corm and the inflorescence; branches (0–)2–5(–9), with (0–)1–3(–6) female heads at the proximal end and (0–)6–9(–16) male heads at the distal end; the upper 1 or rarely 2 branches are often without female heads; the lowest branch often bears a solitary female head without any males, occasionally the solitary head is topped by some males or rarely the branch has male heads only; the terminal




Fig. 18. Fruit set in *Sparganium erectum*; each circle represents a female head, the lower number shows the order of anthesis, the upper number shows the percentage of set fruits (calculated as an average from 85 robust plants growing in the Trent-Mersey Canal, near Derby, Great Britain).

part of the inflorescence (above the last branch) is without female heads and with (3–)5–15(–22) male heads (Fig. 18).

Inflorescence bracts leaf-like below, erect, carinate from base to apex; scale-like above; lowest bract usually exceeding the inflorescence.

Female heads borne on axillary branches, 12–20 mm diameter at anthesis, 15–32 mm diameter in fruit.

Male heads borne above female heads on branches or terminating the main axis, usually remote and distinct at anthesis; before anthesis green with black flecks and ca. 5 mm diameter; at anthesis 10–12 mm diameter.

Female flowers with perianth segments somewhat thickened and dark brown to black at the apex, not translucent, almost entire at apex (Fig. 2a); ovaries with 1, 2 or

rarely 3 carpels but at least half the ovaries are unicarpelate; more than half the styles are undivided; stigmas (1.5–)2–4 mm long, somewhat irregularly bent or coiled.

Male flowers with filaments ca. 5 mm long, anthers 1.2–1.5 mm long; perianth segments thickened and dark brown to black at the apex.

Fruits very variable in form (see subspecies), 4–12 mm long, 2–7.5 mm diameter; peduncle absent or up to 2 mm long, endocarp 3–6(–7) mm long, with sharp to blunt longitudinal ribs (Fig. 4c).

Diagnostic features

Perianth segments thickened and dark brown to black at apex, not translucent and almost entire; inflorescence usually branched; female heads born on axillary branches (very rarely sessile); anthers 1.2–1.5 mm long; less than half the styles are bifid.

Distribution

S. erectum sensu lato is found in Europe from the arctic circle southwards to North Africa extending through temperate Asia southwards to the western Himalaya and eastwards through Japan into northwestern North America (Maps 19–22). It is also in southeastern Australia where it is not certain if it is introduced or native see p. 22.

Ecology

Accounts of the ecology of *S. erectum*, written from different points of view, have been published by Loew in Kirchner et al. (1908), Cook (1962), and Haslam (1978). The autecology of *S. erectum* is very similar to that of *S. emersum* and when both grow together, which is often the case, *S. erectum* occupies the shallower water or it is confined to the banks.

S. erectum is a common species along waterways and in shallow water around ponds, pools and lakes. However, it rarely forms extensive and continuous stands like so many other reed-swamp plants (i.e. *Phragmites*, *Scirpus*, *Typha*); unlike these plants, it is very tolerant to disturbance and can survive considerable battering. In Britain the optimum conditions for growth and flower production (Cook, 1962) are a loose, medium-grained, silty substrate in about 10–20 cm of unshaded, standing or slowly flowing water. In swiftly flowing water the leaves are bent over and there is poor vegetative growth and flowers rarely develop. It can grow in water up to about 2 m deep but again shows poor growth and rarely flowers. In plants stranded on land the immediate response is to remain small (less than 50 cm tall) and develop numerous stolons.

Solitary plants develop leaves which spread fan-like over the water surface. Plants growing in dense stands have almost vertical leaves. The leaves are triangular in transverse section with palisade parenchyma and stomata on all three sides. The spacial geometry of the leaves (see also Kaul, 1973, for *S. eurycarpum*) ensures a high assimilative area. Dykyjová & Ondok (1973) measured a total dry biomass, including below ground organs, exceeding 2500 g/m² in Czechoslovakia; an extremely high level for a component of a natural community in the temperate zone. In spite of this *S. erectum* is a poor competitor and in relatively stable habitats may be overgrown by other erect, amphibious or marginal plants such as, *Glyceria*, *Typha*, or *Phragmites*. In standing water it can invade and replace smaller aquatic plants such as, *Berula erecta*, *Elodea canadensis*, *Ranunculus* spp. or *Rorippa nasturtium-aquaticum*.

S. erectum shows an enormous tolerance and is found in acid, almost oligotrophic waters to brackish or highly polluted water subjected to sewage effluents. In terms of growth and flower production the optimum is mesophytic to eutrophic water somewhat on the alkaline side (pH 7.5–8) in situations that are regularly disturbed (flooding, clearing, dredging, fluctuating water levels, etc). The loose substratum and frequently dirty eutrophic water that the adult plant thrives in are the worst conditions for the juvenile plant. Perhaps reproduction by seed is important in the colonization of newly dug pools and canals where the water is clear and there is no great algal growth. *S. erectum* is effectively spread by detached stolons.

In sluggish streams, canals and in slowly flowing rivers it accumulates silt which may lead to choking. It has been reported to be a serious weed at times in parts of England and the Netherlands. Importation, transportation, and cultivation of *S. erectum* is banned by the Federal Noxious Weed Act of the USA and by the State of Florida; nobody seems to have noticed that it is native in western North America, also it is most unlikely to be able to grow in Florida.

No ecological differences seem to exist between the various subspecies. Cook (1961a) observed four subspecies growing together in the Trent and Mersey canal south of Derby, England. Despite their sympatric distribution, intermediate fruit-shapes were not found.

Notes

S. erectum shows considerable polymorphism in the form and size of the fruits. Polymorphic diaspores are found in several patristically unrelated aquatic plants such as, *Blyxa*, *Ceratophyllum*, *Eclipta*, *Ludwigia* and *Trapa*, see Cook (1987). These examples, like *S. erectum*, show a diaspore polymorphism that is not correlated with vegetative or other floral characteristics. The selective advantage of this polymorphism is difficult to interpret in terms of dispersal function. The taxonomic treatment of these polymorphisms also presents difficulties. In *S. erectum* there are five reasonably distinguishable units which have been variously recognised at ranks from form to species. We find it impracticable to use the rank of species because plants can only be determined with ripe fruits. As each taxon shows a distinct distribution pattern with some allopatry we prefer to adopt the rank of subspecies, which at least allows determination to the rank of species without ripe fruit.

Despite differences in size between fruits of the subspecies of *S. erectum*, our studies show that size is strongly affected by external factors and that size alone may therefore be unreliable as a means of identifying the subspecies.

Plants of *S. erectum* subsp. *neglectum* which had been poorly pollinated (due to deliberate emasculation) were gathered at fruit-set from a pool at the University of Zurich. The number of set fruits in each head was counted and the width and weight of the fruits was measured. These results are presented in Fig. 19. They show a negative correlation between both size and weight of the fruits and the number of fruits per head. Similar size/number tradeoffs have been described, for example, by Galen & Weger (1986).

Ignoring the relative benefits of developing few but large or many but small fruits it was clear to us that *S. erectum* subsp. *neglectum* remained identifiable. From cultivation experiments and other studies we are convinced that the subspecies of *S. erectum* are genetically distinct units and not simply a manifestation of phenotypic plasticity. It is however impossible to distinguish the fruits using size alone; colour and form are the most important characteristics.




Fig. 19. Fruit size (above) and fruit weight (below) in *Sparganium erectum* subsp. *neglectum* from the University of Zürich campus plotted against the number of fruits in each head (the reduced fruit numbers were manipulated by selective pollination).

There are numerous reasons for reduced seed-set in *Sparganium*. Limitations in resources available for maternal function appear to play a role in the number of heads that develop but only influence the number of mature fruits in each head in conditions of stress. Another factor is pollination efficiency; the stigmas remain receptive for, at least, two to three days and pollen is shed over a period of weeks and effectively dispersed by wind. All plants tested were selfcompatible. Nevertheless, there are differences in fecundity between heads depending on their position, see Fig. 18. The lowest head is receptive before the pollen is shed and it is also partly covered by the base of the bract; it frequently shows low fecundity. On all branches the lowest head is usually partially covered by the bract and shows consequently lower seed-set. The uppermost head of the total inflorescence often shows a lower fecundity than its immediate neighbours; this is because it flowers late at a time when most male flowers have shed their pollen and those that are at anthesis are remote at the distal end of the inflorescence.

Some plants are partly sterile and show reduced seed-set on all female heads. The cause of this sterility is unknown but plants marked in nature and plants transferred to the botanic gardens have shown that semi-sterility persists from year to year. This sterility has little to do with pollination efficiency. We suspect it results from hybridization between different subspecies or races of *S. erectum*.

Čelakovský (1896) and his son (Čelakovský, 1899) were convinced that it is possible to distinguish the subspecies of *S. erectum* in Czechoslovakia on the basis of their leaf anatomy. Belavskaja (1984) made similar claims for plants in Russia. We have examined leaf sections of plants growing in different conditions. There is considerable plasticity and we are not convinced that it is possible to distinguish the subspecies on leaf or any other vegetative characteristics.

Hybrid

7×13. *S. emersum* × *S. erectum* sensu lato

= *S. × aschersonianum*, see p. 21

= *S. englerianum*, see p. 24

Experimental crosses have been attempted but the hybrid seems to be inviable. We have no evidence that this hybrid exists, see p. 21 and p. 24.

13. A. ***Sparganium erectum* L. subsp. *erectum***, Species Plantarum 2: 971. 1753. Type: the illustration named *S. ramosum* in L'Obel, Plantarum Seu Stirpium Historia 41. 1576; see Cook (1985) for full discussion.
- = *S. erectum* var. *angustifolium* Warnstorf, Verh. Bot. Verein Brandenburg 37; L [sic]. 2 April 1896 = *S. ramosum* subsp. *polyedrum* subvar. *angustifolium* (Warnstorf) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 293. 15 June 1897 = *S. ramosum* subsp. *polyedrum* var. β *angustifolium* (Warnstorf) Graebner in Engler, Pflanzenreich 2(IV.10): 14. Sept. 1900. Type: Germany, Neu-Ruppin, „im See bei der Wuthenower Fähre“, 1985, Warnstorf (holotype: B, probably destroyed).
- = *S. ramosum* forma *platycarpum* Čelakovský, Österr. Bot. Zeitschrift 46: 423. December 1896 = *S. ramosum* subsp. *polyedrum* subvar. *platycarpum* (Čelakovský) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 284. 15 June 1897 = *S. ramosum* subsp. *polyedrum* var. ϵ *platycarpum* (Čelakovský) Graebner in Engler, Pflanzenreich 2 (IV.10): 14. September 1900. Type: none cited, specimen probably in PR.
- = *S. ramosum* forma *conocarpum* Čelakovský, Österr. Bot. Zeitschrift 46: 423. December 1896 = *S. ramosum* subsp. *polyedrum* subvar. *conocarpum* (Čelakovský) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 284. 15 June 1897 = *S. ramosum* subsp. *polyedrum* var. δ *conocarpum* (Čelakovský) Graebner in Engler, Pflanzenreich 2 (IV.10): 14. September 1900. Type: none cited, specimen probably in PR.
- = *S. ramosum* subsp. *polyedrum* Graebner in Ascherson & Graebner, Syn. Mitteleurop. Flora 1: 283. 15 June 1897. = *S. erectum* subsp. *polyedrum* (Graebner) Schinz & Thellung in Schinz & Keller, Flore Suisse, éd. Française, 1 Partie: Flore d'Excursion, 26. 1909. = *S. ramosum* var. *polyedrum* (Graebner) Holmberg Skand. Flora 1: 79. 1922 = *S. polyedrum* S. W. Juzepczuk, S. V. in Komarov, V. L., Flora USSR. 1: 219. 1934. Type: no specimen cited, based on *S. ramosum* sensu Curtis in W. Curtis, Flora Londinensis 2(fasc. 5): pl. 66 (according to index of fascicle), pl. 194 (according to index of volume 2), pl. 342 (as engraved on the plate, not visible in all copies). April 1787.
- = *S. ramosum* subsp. *polyedrum* subvar. *dolichocarpum* Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 283. 15 June 1897 = *S. ramosum* subsp. *polyedrum* var. γ *dolichocarpum* (Graebner) Graebner in Engler, Pflanzenreich 2 (IV.10): 14. September 1900. Type: Germany, „Westpreußen: Plehnendorf bei Danzig“ (holotype: destroyed in B).




Fig. 20. Fruits, endocarps and diagrammatic sections of fruiting heads of *Sparganium erectum*: a, subsp. *erectum*; b, subsp. *stoloniferum* (scale bar: 1 mm).

- =? *S. draco* F. Hermann, Bull. Soc. Bot. Bulgarie 3: 43. 1929. Type: Bulgaria, "lacu prope Dragoman", specimen not located, not in SOM, perhaps destroyed in B.
- *S. ramosum* Hudson, Flora Anglica, ed. 2, 2: 401. 1778 = *S. erectum* var. δ *ramosum* (Hudson) Laestadius, Wikström's Årsberättelse, 1950, Bihang: 2. 1853 or 1854, nom. illeg. based on *S. erectum* L.
- *S. reyesianum* Sennen, 24 August 1916, *Sennen*, nom. illeg., nom. in schaed.

Fruits cuneate-obpyramidal, (4–)6–10(–12) mm long, (3–)4–6(–7) mm wide, with a distinct shoulder between the upper and lower parts (Fig. 20a); lower part obpyramidal, (–)4.5–8(–10) mm long, distinctly 3–6-angled in transverse section, light brown; upper part flattened light brown to dark brown or blackish, matt (not shiny), smooth (not creased when dry), very abruptly contracted into a beak; beaks not more than 2 mm long; bilocular ovaries frequent but less than half the total ovaries; endocarps (3.5–)5.5–6(–7) mm long, 4–5 mm wide (Fig. 20a), reaching the apex of the fruit; pedicels absent or very short; perianth segments not visible between the fruits on mature fruiting heads, 4.5–7.5 mm long.

Distribution

Common through much of central and southern Europe, but largely absent from Scandinavia. In Britain it is common only south of the Wash. It extends eastwards to the Caucasus (Kaspiysk) and south to Turkey (Hakkari) (see Map 19, the map is based on recently seen herbarium specimens and reliable literature records we have not seen sufficient material from southern and western Europe).

Hybrid

7 × 13 A. S. emersum × S. erectum subsp. *erectum*

= *S. × aschersonianum* Haussknecht, Mitteilungen des Thüringischen Botanischen Vereins N.F., 1 (3–4): 84–86. 1893.

This hybrid is frequently cited in the literature. We very much doubt that it exists. Cook (1962) made some experimental crosses in the field in Derbyshire, England; this cross and the reciprocal cross have each been tried several times without success (the control crosses were fertile). *S. emersum* is often found in a zone occupying slightly deeper water than a parallel zone of *S. erectum* along the banks of canals, streams and small rivers. Pollen must frequently be transferred from one species to the other. We have never found plants in nature that could be referred to this hybrid. Herbarium material that has been determined as this hybrid is almost invariably depauperate and often unbranched *S. erectum*. Authentic material of *S. × aschersonianum* is referable to *S. erectum* subsp. *microcarpum* (p. 22). Rothert (1910) was also of the opinion that this hybrid does not exist.

- 13 B. *Sparganium erectum* subsp. *stoloniferum* (F. Hamilton ex Graebner) C. D. K. Cook & M. S. Nicholls, comb. nova = *S. ramosum* subsp. *stoloniferum* F. Hamilton ex Graebner in Engler, Pflanzenreich 2 (IV.10): 14. September 1900 = *S. stoloniferum* F. Hamilton ex Juzepczuk in Komarov, Flora URSS 1: 219. 1934.
Type: India, Wallich 4990 (holotype: destroyed in B; isotype: K).
= *S. angustifolium* R. Brown, non Michaux, var. *latifolium* Bentham, Fl. Austral. 7: 161. 1878. *Type:* Australia, ?Queensland, Balfours Creek, Leichhardt (holotype: K?; isotype MEL).
= *S. greenei* Morong, Bull Torrey Bot. Club 15: 77, t. 79, fig. 3. 1888 = *S. eurycarpum* var. *greenei* (Morong) Graebner in Engler, Pflanzenreich 2 (IV.10): 13. September 1900. *Type:* USA, California, Marin Co., Olema, September 1887, E. L. Greene (holotype: NY).
= *S. erectum* subsp. *mazanderanicum* Ponert, Fol. Geobot. Phytotax. (Praha) 7: 309. 1972. *Type:* Iran, prov. Mazanderan, "inter oppida Gorgan et Behshahr, 50 km ab Gorgan inter viam ferratam et sinum Gorganicum maris Caspici," 11 July 1970, Jiri Ponert 37919/30, (holotype: BATUMI n.v.; isotype, with male flowers, 37919/29: BATUMI n.v.).
- *S. asiaticum* Graebner, Allg. Bot. Zeitschrift, 4 (2): 32. February 1898, nomen nudum, placed in synonymy of *S. ramosum* subsp. *stoloniferum* by Graebner in Engler, Pflanzenreich 2 (IV.10): 14. September 1900. "Type": specimen used as a basis for Fig. 3c in Pflanzenreich, 2 (IV.10): 12. 1900; B.
- *S. carinatum* Falconer, Proc. Linn. Soc. (London) 1: 18. 1839, never published, nomen illeg., no type cited but hand written manuscript is in K.

Fruits obpyramidal below and domed to conical above, (5–)6–9(–10) mm long, 2.5–5 mm wide; lower part obpyramidal, 4–7 mm long, distinctly 3–6-angled, light

brown; upper part domed to conical with a slight constriction below the shoulder (Fig. 20b), light brown to straw-coloured, shiny, usually with irregular longitudinal creases, tapering rather abruptly into a beak; beak 2–3(–4) mm long; bilocular ovaries common but not more than half the total; endocarps 5–6 mm long, up to 4 mm wide (Fig. 20b), not reaching the apex of the fruit; pedicels absent or very short; perianth segments not visible between the fruits on mature fruiting heads, 4–6 mm long.

Distribution

From the Caucasus and Afghanistan eastwards through north-eastern Pakistan to northern India and Lhasa, Tibet (Map 21). It is recorded from southern China and from west of Peking and occurs through much of Japan except for the extreme north (it is apparently absent from Hokkaido) and also occurs in S.E. Australia where it may have been introduced. It extends into western North America, from California to British Columbia (see Map 22).

Ecology

We have little first-hand knowledge on the ecology of subsp. *stoloniferum*; C.D.K.C. has seen it growing around Srinigar, Kashmir, India. It grows under conditions where one would expect to find other species of *S. erectum* in Europe such as ditches and at the edge of artificially maintained canals, where it is common but rarely builds large stands. In shady conditions, the leaves are often more than 2 m long.

Notes

S. erectum subsp. *stoloniferum* is the most widespread subspecies of *S. erectum*. In Australia it has usually been identified with European *S. erectum* and thus considered to be introduced. It is certainly to be included with the Asian/N. American subspecies *stoloniferum* and on this basis might be considered native in Australia. However, material from this region is abnormally uniform in fruit shape which may indicate that the Australian populations may have developed from a single introduction. In western North America it is also very uniform but is unlikely to be a recent introduction.

13. *C. Sparganium erectum* subsp. *microcarpum* (L. M. Neuman) Domin, Preslia 13: 53. 1935 = *S. ramosum* forma *microcarpa* [sic] L. M. Neuman in Hartman, C. J. & Hartman, C., Handbok Skand. Flora, ed. 12, 122. October 1889 = *S. microcarpum* (L. M. Neuman) L. J. Čelakovský, Österr. Bot. Zeitschrift 66: 378, 423, 1896 = *S. ramosum* subsp. *neglectum* proles *microcarpum* (L. M. Neuman) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 281. 15 June 1897 = *S. erectum* subsp. *neglectum* var. *microcarpum* (L. M. Neuman) A. Hayek, Prodr. Flora Pen. Balcan. Repert. Spec. Nov. Regni Veg. (Fedde), Beiheft, 30 (3): 426. 15 August 1932. Type: Sweden, Gotland, "Visby; Mpd. (Medelpad), Klintehamn"; Per Lassen (LD) could find no specimen collected by *Klintehamn*, it probably perished in the great fire of Sundsvall in 1888. However, there is a specimen from Medelpad collected by Neuman in 1888 (lectotype (designated here): LD – Medelpad: Tuna, Vattjom, 1888, Neuman)

Fruits acutely-obpyramidal below and domed above, (5.5–)6–7(–9) mm long, (2–)2.5–3.5(–4) mm wide; upper and lower parts different in form and texture with a distinct shoulder between them; lower part acutely-obpyramidal, 4–6 mm long, 3–6-




Fig. 21. Fruits, endocarps and diagrammatic sections of fruiting heads of *Sparganium erectum*; a, subsp. *microcarpum*; b, subsp. *neglectum*; c, subsp. *oocarpum* (scale bar: 1 mm).

angled in transverse section, the flattened faces sometimes bellied (endocarp pushes through the side); light brown to reddish brown; upper part domed and wider than the lower part (Fig. 21 a), with a slight constriction below the shoulder, brown to black, matt or somewhat shiny near the shoulder, with irregular longitudinal creases or wrinkles at base of beak, tapering rather abruptly into a beak; beak to 2 mm long, dark-brown to black; bilocular ovaries rare; endocarps (2.5–)4–5 mm long, 2.5–3 mm wide, not reaching the apex of the fruit (Fig. 21 a); pedicels to 1.5 mm long; perianth segments visible on mature fruiting heads, 4–6 mm long.

Distribution

Throughout Europe, particularly common in Scandinavia, extending eastwards to European Russia and south around the Sea of Azov and the Caucasus (Tverksaya). Also from Turkey (Erzurum). See Map 20.

Hybrid

13 A × 13 C. *S. erectum* subsp. *erectum* × *S. erectum* subsp. *microcarpum*

This hybrid is recorded by Ostenfeld-Hansen (1897) and is perhaps the same as

S. ramosum var. ζ *substerile* Neuman, Bot. Not. 1897: 128. 1897 = *S. ramosum* subsp. *polyedrum* var. ζ *substerile* (Neuman) Graebner in Engler, Pflanzenreich 2 (IV.10): 14. 1900. The fruit resembles subsp. *microcarpum* but the heads are usually highly sterile, see p. 17.

13 D. *Sparganium erectum* subsp. *neglectum* (W. H. Beeby) Schinz & Thellung in Schinz & Keller, Flore Suisse, éd. Française, 1 Partie: Flore d'Excursion, 26. 1909 = *S. neglectum* W. H. Beeby, J. Bot. (London), 23: 193, 26, t. 258. 1885 (see also op. cit., 24: 142, 377. 1886) = *S. ramosum* subsp. *neglectum* (W. H. Beeby) L. M. Neuman in Harman, C. J. & Hartman, C., Handbok Skand. Flora, ed. 12, 112. October 1889 or Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 281. 15 June 1897 = *S. erectum* var. β *neglectum* (W. H. Beeby) Richter, Pl. Europ., 1: 10. 1890. Type: England, Surrey, "Albury ponds near Guildford", W. H. Beeby (lectotype: K; isolectotype E).

Fruits fusiform to ellipsoidal, (6-)7-9(-10) mm long, 2-3.5(-5) mm wide, upper and lower parts essentially alike in form, colour and texture, without a distinct shoulder, uniform light brown to straw coloured; lower part 3-5 mm long, about half the fruit length, hardly angled (almost circular in transverse section, Fig. 21 b); upper part smooth (without wrinkles) gradually tapering into a beak; beak (2-)2.5-3.5 mm long, sometimes dark-brown; bilocular ovaries extremely rare; endocarps (4-)5-6 mm long, up to 3.5 mm wide (Fig. 21 b), not reaching the apex of the fruit; pedicels absent or very short; perianth segments visible between the fruits on mature fruiting heads, 5-6 mm long.

Distribution

Throughout much of Europe extending northwards to about 58°N in Sweden and southwards to the northern coast of Africa (Morocco). Also recorded from European Russia and south into Greece and Turkey (Paphlagonia). See Map 19. The published records from Asia Minor are mostly referable to subspecies *stoloniferum*. We have not seen sufficient material from southern and western Europe.

Hybrids

7 × 13 D. *S. emersum* × *S. erectum* subsp. *neglectum*

= *S. × englerianum* Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 287. 15 June 1897. Type: Germany. "Pretzsch", "Berlin: Botanischer Garten" (lectotype: destroyed in B).

We are not convinced that this hybrid exists. Authentic herbarium material refers to depauperate *S. erectum* subsp. *neglectum*.

13 C × 13 D. *S. erectum* subsp. *microcarpum* × *S. erectum* subsp. *neglectum*

This hybrid is probably the same as *S. ramosum* var. ζ *substerile* Neuman, Bot. Not. 1897: 128. 1897 = *S. ramosum* subsp. *polyedrum* var. ζ *substerile* (Neuman) Graebner in Engler, Pflanzenreich 2 (IV.10): 14. 1900. It resembles subsp. *microcarpum* but the fruits are somewhat larger and elongated and the heads are partly sterile. It is occasionally found growing with the parents.

13 A × 13 D. *S. erectum* subsp. *erectum* × *S. erectum* subsp. *neglectum*

= *S. × tardivum*, Topa in Săvulescu, T. Flora Republicii Socialiste România 11: 849 (see

also p. 106) 1966. Type: Romania, "Reg. Iași, Horlești, propr. oppidum Iași, legit. *M. Răvărut*" (holotype: I or possibly IAGB, n.v.).

Several authors including Cook (in Stace, 1975) have suggested that *S. erectum* subsp. *oocarpum* is this hybrid, while others such as Čelakovský (1896) were convinced that it is not a hybrid. We note that subsp. *oocarpum* is not fully fertile, it well may be of hybrid origin but is most unlikely to be a simple cross between subsp. *erectum* and subsp. *neglectum* (for further discussion see p. 17).

- 13 E. *Sparganium erectum* subsp. *oocarpum* (L. J. Čelakovský) Domin, Preslia 13: 53. 1935 = *S. neglectum* var. *oocarpum* L. J. Čelakovský, Österr. Bot. Zeitschrift 46: 425. December 1896 = *S. ramosum* subsp. *neglectum* proles *oocarpum* (L. J. Čelakovský) Graebner in Ascherson & Graebner, Synopsis Mitteleurop. Flora 1: 282. 15 June 1897 = *S. neglectum* subsp. *oocarpum* (L. J. Čelakovský) C. Ostenfeld-Hansen, Bot. Tidsskr. 21 (1): Meddelesler V. 6 July 1897 = *S. oocarpum* (L. J. Čelakovský) Fritsch, Exkursionsflora Österr., ed. 2, 28. 1909 = *S. erectum* subsp. *neglectum* var. *oocarpum* (L. J. Čelakovský) Hayek, Prodr. Flora Pen. Balcan. Repert. Spec. Nov. Regni Veg. (Fedde), Beiheft, 30 (3): 426. 15 August 1932. Type: Czechoslovakia, several localities given (lectotype: PR to be chosen).
 = *S. × tardivum*, Topa in Săvulescu. T. Flora Republicii Socialiste România 11: 849 (see also p. 106) 1966. Type: Romania, "Reg. Iași, Horlești, propr. oppidum Iași, legit. *M. Răvărut*" (holotype: I or possible IAGB, n.v.).

Fruits widely ovoid to almost spherical, 5–8 mm long, 4–7 mm wide, uniform light to medium brown and shiny, shoulder between upper and lower parts indistinct; lower part 2.5–5 mm long, almost circular in transverse section; upper part tapering into a beak and sometimes wrinkled or creased at base of beak but otherwise smooth (Fig. 21 c); beaks up to 2 mm long; bilocular ovaries very rare; endocarps 3.5–4 mm long, up to 4 mm wide; not reaching the apex of the fruit, pedicels absent or very short; perianth segments visible between the fruits on mature fruiting heads, 3–6 mm long (Fig. 21 c).

Distribution

Isolated records through much of Europe, Turkey (Istanbul, Vol Antalya) and Iraq (Map 20).

Notes

Specimens show very poor fertility which has led Cook (in Stace, 1975) and others to suggest that subsp. *oocarpum* is of hybrid origin. On morphological grounds, the suggested parents are subsp. *erectum* and subsp. *neglectum* but it is unlikely to be a simple cross between these two.

It has also been suggested that the big fat fruits of subsp. *oocarpum* are a consequence of resource allocation (caused, for example, by poor pollination). A plant with few fruits to mature per head might be expected to devote greater resources to the remaining fruits which will then grow fatter than fruits competing more strongly for resources. We have undertaken experiments with subsp. *neglectum* where pollination was artificially limited so that few ovules per head were fertilized. Under these conditions subsp. *neglectum* also produces "fat" fruits and both fruit-width and fruit-weight are strongly correlated with the number of fruits per head (see Fig. 19). Nevertheless, these

fruits are still distinguishable from true *oocarpum* leading us to discount this resource allocation hypothesis as a means of accounting for the origin of this subspecies.

14 A. ***Sparganium eurycarpum* subsp. *eurycarpum*** Engelmann in Gray, Manual Bot. Northern U.S., ed. 2, 430. 1856. Type: USA, "from New England and Pennsylvania northward and westward", Engelmann (lectotype to be chosen from material at MO; possible isotype: G).

= *S. californicum* Greene, Bull. Calif. Acad., 1 (3): 11. 1884. Type: USA, California, Calistoga, E. L. Greene (holotype: ND-G; isotype: ?NY).

Robust erect, emergent or rarely floating or submerged herbs. Stolons up to ca. 30 cm long, ca. 5 mm diameter. Corms up to 4.5 cm diameter, when mature hard and woody.

Basal leaves obliquely erect and fan-like, 50–100(–260) cm long, 6–20 mm wide, carinate at base but often becoming flattened towards the apex, with 3–6 layers of gas chambers.

Plants in very deep or swiftly flowing water may develop submerged or partly floating leaves but usually remain sterile.

Flowering stems erect, 50–150(–260) cm long, nearly always branched with 1–4 sterile nodes between the corm and the inflorescence; branches (0–)2–5, axillary, with 1–2 or rarely more female heads at the proximal end and (0–)6–14 male heads at the distal end or upper branches with male heads only; the upper female head or heads are oc-




Fig. 22. Fruit and endocarp of *Sparganium eurycarpum* subsp. *eurycarpum* (scale bar: 1 mm).

casionally sessile on main axis; the lowest branch usually with both male and female heads.

Inflorescence bracts leaf-like below, erect, carinate at base; scale-like above; lowest bract usually not exceeding inflorescence at late anthesis.

Female heads borne on axillary branches, peduncles or sessile on main axis, 2–3.5 mm diameter in fruit (excluding beaks).

Male heads borne above female heads on branches or terminating the main axis, usually remote and distinct at anthesis; before anthesis green with black flecks and ca. 5 mm diameter; at anthesis ca. 12 mm diameter.

Female flowers with perianth segments somewhat thickened and dark brown to black at the apex, not translucent almost entire at apex; ovaries at least half with 2 carpels the majority of the rest with 1 carpel and some with 3 carpels; at least half the stigmas bifid; stigmas 2–3.5 mm long, usually curved.

Male flowers with filaments ca. 6 mm long; anthers 1–1.5 mm long, perianth segments somewhat thickened and dark-brown to black at apex.

Fruits cuneate-obpyramidal, 6–10 mm long, 6–8 mm wide with a distinct shoulder between upper and lower parts (Fig. 22); lower part obpyramidal, 5–9 mm long, distinctly 3–6 angled in transverse section, light brown; upper part flattened, straw-coloured to brown, sometimes sootyblack at base of beak, mostly smooth (not creased when dry), very abruptly contracted into a beak; beaks 2–4(–4) mm long; endocarps 7–10 mm long, up to 7 mm wide, usually 2-locular, distinctly ribbed with ribs projecting beyond the base (Fig. 22), reaching the apex of the fruit; pedicels absent; perianth segments not visible between the fruits on mature fruiting heads.

Diagnostic features

Like *S. erectum* but at least half the styles bifid (at least half the ovaries have two carpels); fruits sessile, cuneate-obpyramidal, 6–10 mm long, 6–8 mm wide, the top truncate depressed or very shallowly rounded, abruptly tipped with the style, straw coloured to somewhat sooty-black.

Distribution

It occurs across North America, from east to west between 60°N and 35°N, (Map 22).

Ecology

S. eurycarpum is essentially a north American vicariant of the eurasian species *S. erectum*. Both species are ecologically very similar and although we have no experience of *S. eurycarpum* in the field, most of the information presented on pp. 16–18 is probably relevant. It is reported to grow on damp shores and in still or slowly flowing water. It performs best in slightly alkaline conditions but tolerates acid substrates. It is often very abundant although not reported to be a serious weed. Rather surprisingly, it is on the rare and endangered plant list for New Hampshire.

Neely & Davis (1985a) found that fertilization of *S. eurycarpum* with nitrogen and phosphorus increased the net annual above-ground production but the root-rhizome production was not altered. The litter is a poor nutrient sink and Neely & Davis (1985b) noted that litter with an initially high nutrient content decomposed faster than litter with low initial nutrient content.

Cruden & Lyon (1985) showed that in *S. eurycarpum* the dry weight of both male and female heads increased with the number of inflorescence branches, and that dry weight is positively correlated with the number of seeds matured. They interpret these findings in the light of current theories regarding resource allocation patterns in plants.

Although Cruden & Lyon's findings are hardly surprising, we would question their assumption that *S. eurycarpum* is xenogamous, particularly in light of our own work on pollen-flow pattern in *S. erectum* (see p. 18).

- 14 B. ***Sparganium eurycarpum* subsp. *coreanum*** (Léveillé) C. D. K. Cook & M. S. Nicholls comb. nova = *S. coreanum* Léveillé, Repert. Spec. Nov. Regni Veg. (Fedde) 10: 441. 1912 = *S. stoloniferum* var. *coreanum* (Léveillé) Hara fide Ohwi, Flora Japan 119. 1965. Type: S. Korea, Cheju do, "Quelpaert, in stagno Terok", 14 September 1908. Taquet 2150 (holotype: E; isotypes: G, K, TI).
 – *S. macrocarpum* Makino in Matsumura, Indig. Pl. Jap. (2) 1: 24. 1905, nomen nudum = *S. stoloniferum* var. *macrocarpum* (Makino) Hara fide Ohwi, Flora Japan 119. 1965. Type: Japan, Honshu, Chiba Pref., Junsai-numa, September 1893, T. Makino s.n. (holotype: TI).

Like subsp. *eurycarpum* but mature fruiting heads usually smaller, rarely exceeding 20 mm diameter (excluding beaks); individual fruits 5–9(–10) mm long, 5–8 mm wide; the top of the fruit less flat, shallowly rounded or widely pyramidal (Fig. 23) and light




Fig. 23. Fruit and endocarp of *Sparganium eurycarpum* subsp. *coreanum* (scale bar: 1 mm).

to dark brown, not shiny, abruptly contracted into a 2–4 mm long beak; endocarps 6–7 mm long, 4–5 mm wide with longitudinal ribs extending beyond the base (Fig. 23).

Distribution

Korea to northern and central Japan (Honshu). Also from Manchuria (see Map 21).

Notes

We have seen few adequate specimens but are, nonetheless, convinced that the affinities of this taxon lie with *S. eurycarpum* rather than *S. erectum*. The fruits are very large and more than half are bilocular with bifid styles.

Uncertain species

- Sparganium affine* forma *abbreviata* Meyerholz. Verh. Bot. Verein Brandenburg 34: 26. 1893, nomen nudum, no authentic specimens found.
- Sparganium alpinum* D. Don ex G. Don in Loudon, Hortus Britannicus, 375. 1830, nomen nudum, no authentic specimens found.
- Sparganium androcladum* var. ζ *microcarpum* Graebner in Engler, Pflanzenreich 2 (IV.10): 15. 1900, diagnosis inadequate, no authentic specimens found.
- Sparganium axilare* Rafinesque, Autikon Botanikon, 198. 1840, description inadequate, no authentic specimens found, see Cook (1985) for details.
- Sparganium emersum* forma *angustifolium* (Morong) [sic] Soó, Acta Bot. Acad. Sci. Hung. 17 (1–2): 124. 1971 publ. 1972, perhaps a misprint.
- Sparganium ligulare* Rafinesque, Autikon Botanikon, 198. 1840, description inadequate, no authentic specimens found, see Cook (1985) for details.
- Sparganium simplex* var. *subnatans* E. M. Fries, Bot. Not. 1868: 71. 1868, nomen nudum, no authentic specimens found.
- Sparganium subspinosa* Auct., Just's Bot. Jahresber. 21 (2): 670. 1893, name appears in index, probably a misprint or mistake.

References

- Beal E. O. 1960. *Sparganium* (Sparganiaceae) in the southeastern United States. Brittonia 12: 176–181.
- Belavskaja A. P. 1984. A contribution to the morphology of fruits of the genus *Sparganium* (Typhaceae) in the flora of the USSR. Botaniceskij Žurnal Akad. Nauk. SSSR (Leningrad) 69 (12): 1662–1668.
- Čelakovský L. J. 1896. Über die ramosen Sparganien Böhmens. Österreich. Bot. Zeitschr. 46: 377–381, 46: 421–433.
- Čelakovský L. 1899. Anatomický rozdíly v Pistech ramoních sparganii. Věstn. Král. Čes. Společ. Nauk. Cl. Math.-Natur. (Praha) 1899 (5): 1–11.
- Chen Yao-Dong 1981. A study on Chinese *Sparganium* (in Chinese). Acta Phytotaxonomica Sinica 19 (1): 41–57.
- Clausen R. T. 1937. A new variety of *Sparganium americanum*. Rhodora 39: 188–190.
- Cook C. D. K. 1961. *Sparganium* in Britain. Watsonia 5: 1–10.
- Cook C. D. K. 1962. Biological Flora of the British Isles, No. 82. *Sparganium erectum* L. J. Ecology 50: 247–255.

- Cook C. D. K. 1985. *Sparganium*: some old names and their types. Bot. Jahrb. Syst. 107: 269–276.
- Cook C. D. K. 1987. Dispersion in aquatic and amphibious vascular plants. In Crawford (Ed.) Plant Life in Aquatic and Amphibious Habitats. British Ecological Society Special Symposium Blackwell, Oxford (in press).
- Cruden R. W. & Lyon D. L. 1985. Pattern of biomass allocation to male and female functions in plants with different mating systems. Oecologia (Berlin) 66: 299–306.
- Dykyjová D. & Ondok J. P. 1973. Biometry and the productive stand structure of coenoses of *Sparganium erectum*. L. Preslia (Praha) 45: 19–30.
- Fernald M. L. 1922. Notes on *Sparganium*. Rhodora 24(277): 26–34.
- Fernald M. L. & Eames A. J. 1907. Preliminary lists of New England Plants, XX. Spanganiceae. Rhodora 8 (85): 86–90.
- Galen C. & Weger H. G. 1986. Re-evaluating the significance of the correlations between seed number and size: evidence from a natural population of the lily, *Clintonia borealis*. Amer. J. Bot. 73: 346–352.
- Haslam S. M. 1978. River plants. Cambridge University Press, Cambridge. pp. 1–396.
- Kaul R. B. 1973. Development of foliar diaphragms in *Sparganium eurycarpum*. Amer. J. Bot. 60: 944–949.
- Kirchner O., Loew E. & Schroeter C. 1908. Lebensgeschichte der Blütenpflanzen Mitteleuropas 1 (1): 374–394. Eugen Ulmer, Stuttgart.
- Mohr C. 1901. Plant life of Alabama. Contrib. U.S. Nat. Herbarium 6: 327.
- Morong T. 1888. Studies in the Typhaceae. II. *Sparganium*. Bull. Torrey Bot. Club 15: 73–81.
- Neely R. K. & Davis C. B. 1985 a. Nitrogen and phosphorous fertilization of *Sparganium eurycarpum* Engelm. and *Typha glauca* Godr. stands. I. Emergent plant production. Aquatic Bot. 22: 347–361.
- Neely R. K. & Davis C. B. 1985 b. Nitrogen and phosphorous fertilization of *Sparganium eurycarpum* Engelm. and *Typha glauca* Godr. stands. II. Emergent plant decomposition. Aquatic Bot. 22: 363–375.
- Nuttall T. 1818. The genera of North American plants. Facsimile ed.: 1971. Hafner Publ. Co. New York. Classica botanica americana vol. 7.
- Ostenfeld-Hansen C. 1897. De i Danmark voxende ramøse *Sparganium*-Arten. Bot. Tidskrift Bot. For. Kjøbenhavn 21 (1): V–IX.
- Rothert W. 1910. Übersicht der Sparganien des Russischen Reiches (zgleich Europa's). Acta Horti Bot. Univ. Imper. Jurjevensis 11: 11–32.
- Rydberg P. A. 1909. Sparganiaceae in North American Flora (New York Botanical Gardens) 17: 5–10.
- Stace C. A. 1975. Hybridization and the Flora of the British Isles. Academic Press, London, New York, San Francisco. pp. 1–626.

Index

Index to names, combinations and synonyms used in this study. **Bold**: accepted names; *italic*: synonyms or illegitimate names. “1”: indicates part 1, Bot. Helv. 96: 1–56 (1986); “2”: in this part.

Isoetes lacustris var. *fluitans* Döll 1:238

Sparganium

<i>acaule</i> (Beeby) Rydb.	1:257
<i>affine</i> Schnitzl.	1:238
<i>affine</i> Schnitzl subsp. <i>borderi</i> (Focke) Weberb. ex Graeb.	1:238
<i>affine</i> Schnitzl subsp. <i>borderi</i> var. <i>deminutum</i> (Neuman) Graeb.	1:239
<i>affine</i> Schnitzl subsp. <i>borderi</i> var. <i>microcephalum</i> (Neuman) Graeb.	1:239
<i>affine</i> Schnitzl var. <i>deminutum</i> Neuman	1:239
<i>affine</i> Schnitzl var. <i>microcephalum</i> Neuman	1:239


<i>affine</i> Schnitzl var. <i>zosteraefolium</i> [sic] Neuman	1:238
<i>affine</i> Schnitzl subvar. <i>zosterifolium</i> (Neuman) Graeb.	1:238
<i>affine</i> Schnitzl forma <i>abbreviata</i> Meyerh.	2:29
<i>alpinum</i> D. Don ex G. Don	2:29
<i>americanum</i> Nuttall	2:10
<i>americanum</i> var. <i>androcladum</i> (Engelm.) Fern. & Eames	2:13
<i>americanum</i> var. <i>rigidum</i> Clausen	2:10
<i>androcladum</i> (Engelmann) Morong	2:12
<i>androcladum</i> (Engelmann) var. <i>fluctuans</i> Morong	1:247
<i>androcladum</i> (Engelmann) var. <i>microcarpum</i> Graeb.	2:29
<i>angustifolium</i> Mich.	1:238
<i>angustifolium</i> subsp. <i>emersum</i> (Rehm.) Brayshaw	1:249
<i>angustifolium</i> subsp. <i>emersum</i> var. <i>chlorocarpum</i> (Rydb.) Brayshaw	1:250
<i>angustifolium</i> subsp. <i>emersum</i> var. <i>multipedunculatum</i> (Morong) Brayshaw	1:238
<i>angustifolium</i> R. Br. var. <i>latifolium</i> Bentham	2:21
<i>angustifolium</i> × <i>emersum</i>	1:244, 256
<i>angustifolium</i> × <i>glomeratum</i>	1:244
<i>angustifolium</i> × <i>gramineum</i>	1:247, 248, 257
<i>angustifolium</i> × <i>hyperboreum</i>	1:242
<i>angustifolium</i> × <i>natans</i>	1:242
<i>antipodum</i> Graeb.	2:4
<i>aschersonianum</i> Hausskn.	2:21
<i>asiatium</i> Graeb.	2:21
<i>axilare</i> Rafin.	2:29
<i>balticum</i> Dorofeev	1:219
<i>barrerae</i> Sennen	2:14
<i>boreale</i> Least. ex Beurl.	1:255
<i>borderi</i> Focke	1:238
<i>californicum</i> Greene	2:26
<i>carinatum</i> Falconer	2:21
<i>chlorocarpum</i> Rydb.	1:250
<i>chlorocarpum</i> Rydb. var. <i>acaule</i> (Beeby) Fern.	1:257
<i>chlorocarpum</i> Rydb. forma <i>acaule</i> (Beeby) Voss	1:257
<i>confertum</i> Y.-D. Chen	2:2
<i>coreanum</i> Lév.	2:28
<i>costatum</i> Dorofeev	1:219
<i>diversifolium</i> Graeb.	1:250, 255
<i>diversifolium</i> Graeb. var. <i>acaule</i> (Beeby) Fern. & Eames	1:257
<i>diversifolium</i> Graeb. proles <i>nanum</i> Graeb.	1:257
<i>diversifolium</i> Graeb. proles <i>wirtgeniorum</i> Graeb.	1:250
<i>draco</i> Hermann	2:20
<i>emersum</i> Rehmann	1:249
<i>emersum</i> subsp. <i>acaule</i> (Beeby) Cook & Nicholls	1:257
<i>emersum</i> subsp. <i>simplex</i> (Hudson) Soó	1:251
<i>emersum</i> var. <i>angustifolium</i> (Mich.) Taylor & McBride	1:238
<i>emersum</i> var. <i>multipedunculatum</i> (Morong) Reveal	1:238
<i>emersum</i> forma <i>angustifolium</i> (Morong) Soó	2:29
<i>emersum</i> forma <i>natans</i> (Glück ex) Soó	1:251
<i>emersum</i> forma <i>simile</i> (Meinsh.) Soó	1:250
<i>emersum</i> forma <i>splendens</i> (Meinsh.) Soó	1:250

<i>emersum</i> forma <i>submersum</i> (Glück ex) Soó	1:251
<i>emersum</i> forma <i>subvaginatum</i> (Meinsh.) Soó	1:239, 250
<i>emersum</i> × <i>angustifolium</i>	1:255
<i>emersum</i> × <i>erectum</i> s.l.	2:19
<i>emersum</i> × <i>erectum</i> subsp. <i>erectum</i>	2:21
<i>emersum</i> × <i>erectum</i> subsp. <i>neglectum</i>	2:24
<i>emersum</i> × <i>glomeratum</i>	1:257
<i>emersum</i> × <i>gramineum</i>	1:257
<i>emersum</i> × <i>hyperboreum</i>	1:255
<i>emersum</i> × <i>natans</i>	1:255
<i>englerianum</i> Graeb.	2:19, 24
<i>erectum</i> L. sensu lato	2:14
<i>erectum</i> subsp. <i>erectum</i> L.	2:19
<i>erectum</i> subsp. <i>mazanderanicum</i> Ponert	2:21
<i>erectum</i> subsp. <i>microcarpum</i> (Neuman) Domin	2:22
<i>erectum</i> subsp. <i>neglectum</i> (Beeby) Schinz & Thell.	2:24
<i>erectum</i> subsp. <i>neglectum</i> var. <i>microcarpum</i> (Neuman) Hayek	2:22
<i>erectum</i> subsp. <i>neglectum</i> var. <i>oocarpum</i> (Čelak.) Hayek	2:25
<i>erectum</i> subsp. <i>oocarpum</i> (Čelak.) Domin	2:25
<i>erectum</i> subsp. <i>polyedrum</i> (Graeb.) Schinz & Thell.	2:19
<i>erectum</i> subsp. <i>stoloniferum</i> (F. Ham. ex Graeb.) Cook & Nicholls	2:21
<i>erectum</i> var. <i>angustifolium</i> Warnst.	2:19
<i>erectum</i> var. <i>boreale</i> Least.	1:251, 255
<i>erectum</i> var. <i>glomeratum</i> Laest.	1:243
<i>erectum</i> var. <i>neglectum</i> (Beeby) Richter	2:24
<i>erectum</i> var. <i>non-ramosum</i> L.	1:249
<i>erectum</i> var. <i>ramosum</i> Laest.	2:20
<i>erectum</i> var. <i>simplex</i> Hudson) Laest.	1:252
<i>erectum</i> s.l. × <i>emersum</i>	2:19
<i>erectum</i> subsp. <i>erectum</i> × <i>emersum</i>	2:21
<i>erectum</i> subsp. <i>erectum</i> × <i>erectum</i> subsp. <i>microcarpum</i>	2:23
<i>erectum</i> subsp. <i>erectum</i> × <i>erectum</i> subsp. <i>neglectum</i>	2:24
<i>erectum</i> subsp. <i>microcarpum</i> × <i>erectum</i> subsp. <i>erectum</i>	2:23
<i>erectum</i> subsp. <i>microcarpum</i> × <i>erectum</i> subsp. <i>neglectum</i>	2:24
<i>erectum</i> subsp. <i>neglectum</i> × <i>emersum</i>	2:24
<i>erectum</i> subsp. <i>neglectum</i> × <i>erectum</i> subsp. <i>erectum</i>	2:24
<i>erectum</i> subsp. <i>neglectum</i> × <i>erectum</i> subsp. <i>microcarpum</i>	2:24
<i>eurycarpum</i> Engelmann	2:26
<i>eurycarpum</i> subsp. <i>coreanum</i> (Lév.) Cook & Nicholls	2:28
<i>eurycarpum</i> var. <i>greenei</i> (Morong) Graeb.	2:21
<i>fallax</i> Graeb.	2:2
<i>fallax</i> × <i>subglobosum</i>	2:4, 8
<i>flaccidum</i> Meinsh.	1:235
<i>fluctuans</i> (Morong) Robinson	1:247
<i>fluitans</i> (E. M. Fries) E. M. Fries	1:243
<i>friesii</i> Beurling	1:245
<i>glehnii</i> Meinsh.	1:243
<i>glomeratum</i> (Beurl. ex Laest.) Neuman	1:242
<i>glomeratum</i> var. <i>angustifolium</i> Graeb.	1:244
<i>glomeratum</i> × <i>angustifolium</i>	1:244
<i>glomeratum</i> × <i>emersum</i>	1:244


gramineum Georgi	1:244
gramineum × <i>angustifolium</i>	1:247, 257
gramineum × <i>emersum</i>	1:247, 257
<i>greenei</i> Morong	2:21
hyperboreum Laest. ex Beurl.	1:231
hyperboreum var. <i>natans</i> Beurl.	1:231
<i>hyperboreum</i> × <i>angustifolium</i>	1:234
<i>hyperboreum</i> × <i>emersum</i>	1:234
<i>hyperboreum</i> × <i>natans</i>	1:234, 237
japonicum Rothert	2:8
<i>kawakamii</i> Hara	1:239
<i>lanceolatum</i> Georgi	1:245
<i>ligulare</i> Rafin.	2:29
<i>limosum</i> Y.-D. Chen	2:6
<i>longifolium</i> Turcz.	1:257
<i>longissimum</i> (Fries) Fritsch	1:251
<i>lucidum</i> Fern. & Eames	2:13
<i>macrocarpum</i> Makino	2:28
<i>microcarpum</i> (Neuman) Čelak.	2:22
<i>microcarpum</i> × <i>ramosum</i>	2:23
<i>minimum</i> Ray, Hill, Wallroth, E. M. Fries	1:234
<i>minimum</i> var. <i>flaccidum</i> (Meinsh.) Graeb.	1:235
<i>minimum</i> var. subvar. <i>rostratum</i> (Larsson) Graeb.	1:235
<i>minimum</i> var. subvar. <i>oligocarpon</i> (Ångst.) Graeb.	1:235
<i>minimum</i> var. subvar. subvar. <i>ratis</i> (Meinsh.) Graeb.	1:235
<i>minimum</i> var. subvar. subvar. <i>septentrionale</i> (Meinsh.) Graeb.	1:235
<i>minimum</i> var. subvar. subvar. <i>perpusillum</i> (Meinsh.) Graeb.	1:235
<i>minimum</i> var. subvar. subvar. <i>strictum</i> Luersson	1:235
<i>minimum</i> var. forma <i>rostrata</i> [sic] (Larsson) Neuman	1:235
multiloculare Reid & Chandler	1:219
multipedunculatum (Morong) Rydb.	1:238
natans L.	1:234
<i>natans</i> L. var. <i>angustifolium</i> (Mich.) Pursh	1:238
<i>natans</i> L. var. <i>minimum</i> Hartman	1:234
<i>natans</i> L. var. <i>subdecumbens</i> Laest.	1:235
<i>natans</i> L. var. <i>suberectum</i> Beurl. ex Laest.	1:235
<i>natans</i> L. var. <i>submuticum</i> Hartman	1:231
<i>natans</i> L. forma <i>gracilis</i> Neuman	1:250
<i>natans</i> × <i>angustifolium</i>	1:237
<i>natans</i> × <i>emersum</i>	1:237, 255
<i>natans</i> × <i>hyperboreum</i>	1:237
<i>neglectum</i> Beeby	2:24
<i>neglectum</i> subsp. <i>oocarpum</i> (Čelak.) Ostenfeld-Hansen	2:25
<i>neglectum</i> var. <i>oocarpum</i> Čelak.	2:25
<i>nipponicum</i> Makino	2:6
<i>oligocarpon</i> Ångst.	1:238
<i>oocarpum</i> (Čelak.) Fritsch	2:25
<i>perpusillum</i> Meinsh.	1:235
<i>polyedrum</i> (Graeb.) Juz.	2:19
<i>pubescens</i> Poiret	1:216

<i>ramosum</i> Hudson	2:20
<i>ramosum</i> subsp. <i>neglectum</i> (Beeby) Neuman	2:24
<i>ramosum</i> subsp. <i>neglectum</i> var. <i>microcarpum</i> (Neuman) Hayek	2:22
<i>ramosum</i> subsp. <i>neglectum</i> var. <i>microcarpum</i> proles <i>microcarpum</i> (Neuman) Graeb.	2:22
<i>ramosum</i> subsp. <i>neglectum</i> var. <i>microcarpum</i> proles <i>oocarpum</i> (Čelak.) Graeb.	2:25
<i>ramosum</i> subsp. <i>polyedrum</i> Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> var. <i>angustifolium</i> (Warnst.) Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> var. <i>conocarpum</i> (Čelak.) Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> var. <i>dolichocarpum</i> (Graeb.) Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> var. <i>platycarpum</i> (Čelak.) Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> var. <i>substerile</i> (Neuman) Graeb.	2:24
<i>ramosum</i> subsp. <i>polyedrum</i> subvar. <i>angustifolium</i> Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> subvar. <i>conocarpum</i> (Čelak.) Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> subvar. <i>dolichocarpum</i> Graeb.	2:19
<i>ramosum</i> subsp. <i>polyedrum</i> subvar. <i>platycarpum</i> (Čelak.) Graeb.	2:19
<i>ramosum</i> subsp. <i>stoloniferum</i> F. Ham. ex Graeb.	2:21
<i>ramosum</i> var. <i>polyedrum</i> (Graeb.) Holmb.	2:19
<i>ramosum</i> var. <i>substerile</i> Neuman	2:24
<i>ramosum</i> forma <i>conocarpum</i> Čelak.	2:19
<i>ramosum</i> forma <i>microcarpa</i> [sic] Neuman	2:22
<i>ramosum</i> forma <i>platycarpum</i> Čelak.	2:19
<i>ramosum</i> forma <i>simplicior</i> Rothert	2:14
<i>ramosum</i> × <i>simplex</i>	2:19
<i>ramosum</i> subsp. <i>neglectum</i> × <i>simplex</i>	2:24
<i>ramosum</i> subsp. <i>polyedrum</i> × <i>simplex</i>	2:21
<i>ratis</i> Meinh.	1:235
<i>reyesianum</i> Sennen	2:20
<i>rostratum</i> Larsson	1:235
<i>septentrionale</i> Meinh.	1:235
<i>simile</i> Meinh.	1:250
<i>simplex</i> Hudson	1:251
<i>simplex</i> var. <i>acaule</i> Beeby	1:257
<i>simplex</i> var. <i>americanum</i> (Engelm.) Engler	2:10
<i>simplex</i> var. <i>androcladum</i> Engelm.	2:12
<i>simplex</i> var. <i>androgyna</i> [sic] Meinh.	2:13
<i>simplex</i> var. <i>angustifolium</i> Beckmann	1:251
<i>simplex</i> var. <i>angustifolium</i> (Michaux) Engelm.	1:239
<i>simplex</i> var. <i>angustifolium</i> subvar. <i>gracile</i> [sic] (Meinh.) Graeb.	1:250
<i>simplex</i> var. <i>angustifolium</i> subvar. <i>subvaginatum</i> (Meinh.) Graeb.	1:239, 250
<i>simplex</i> var. <i>fluitans</i> Engelm.	1:247
<i>simplex</i> var. <i>fluitans</i> E. M. Fries	1:242
<i>simplex</i> var. <i>fluitans</i> God. & Gren.	1:250
<i>simplex</i> var. <i>glomeratum</i> Laest.	1:242
<i>simplex</i> var. <i>gracile</i> Meinh.	1:250
<i>simplex</i> var. <i>longissimum</i> (E. M. Fries) Neuman	1:251
<i>simplex</i> var. <i>multipedunculata</i> [sic] Morong	1:238
<i>simplex</i> var. <i>nuttallii</i> Engelm.	2:10
<i>simplex</i> var. <i>simile</i> (Meinh.) Graeb.	1:250
<i>simplex</i> var. <i>splendens</i> (Meinh.) Graeb.	1:250
<i>simplex</i> var. <i>splendens</i> subvar. <i>simile</i> (Meinh.) Graeb.	1:250
<i>simplex</i> var. <i>subnatans</i> E. M. Fries	2:29
<i>simplex</i> var. <i>subvaginatum</i> (Meinh.) Graeb.	1:239
<i>simplex</i> var. proles <i>longissimum</i> (E. M. Fries) Graeb.	1:251


<i>simplex</i> var. <i>proles</i> var. <i>emersum</i> (Rehm.) Graeb.	1:249
<i>simplex</i> var. <i>proles</i> var. <i>inundatum</i> (Graeb.) Graeb.	1:251
<i>simplex</i> var. <i>proles</i> var. <i>multipedunculatum</i> (Morong) Graeb.	1:238
<i>simplex</i> var. <i>proles</i> var. subvar. <i>emersum</i> (Rehm.) Graeb.	1:249
<i>simplex</i> var. <i>proles</i> var. subvar. <i>inundatum</i> Graeb.	1:251
sobolevii Dorofeev	1:219
<i>speirocephalum</i> Neuman	1:257
<i>splendens</i> Meinh.	1:250
<i>stenophyllum</i> Maxim. ex Meinh.	2:4
<i>stolonifeum</i> F. Ham. ex Juz.	2:21
<i>stolonifeum</i> var. <i>coreanum</i> (Lév.) Hara	2:28
<i>stolonifeum</i> var. <i>macrocarpum</i> (Makino) Hara	2:28
subglobosum Morong	2:4
<i>subglobosum</i> × <i>fallax</i>	2:4, 8
<i>submuticum</i> (Hartman) Neuman	1:231
<i>submuticum</i> forma <i>platyphylla</i> [sic] Neuman	1:231
<i>subvaginatum</i> Meinh.	1:239, 250
<i>subspinosa</i> [sic] Just	2:29
<i>tardivum</i> Topa	2:21, 25
<i>tenuifolium</i> Poiret	1:216
<i>trifidum</i> Poiret	1:216
<i>vaginatum</i> Larsson	1:238
<i>williamsii</i> Rydb.	1:231
<i>wirtgeniorum</i> (Graeb.) Rouy	1:250
<i>yamatense</i> Y.-D. Chen	2:2
<i>yunnanense</i> Y.-D. Chen	2:2
<i>zetlandicum</i> Druce	1:255




Map 14. *Sparganium fallax*.




Map 15. *Sparganium subglobosum*.




Map 16. *Sparganium japonicum*.




Map 17. *Sparganium americanum*.




Map 18. *Sparganium androcladum*.




Map 19. *Sparganium erectum* subsp. *erectum* (▲), subsp. *neglectum* (●).



Map 20. *Sparganium erectum* subsp. *microcarpum* (●), subsp. *oocarpum* (▲).



Map 21. *Sparganium erectum* subsp. *stoloniferum* (●) and *S. eurycarpum* subsp. *coreanum* (▲) in Asia and Australia (inset).



Map 22. *Sparganium eurycarpum* subsp. *eurycarpum* (●) and *S. erectum* subsp. *stoloniferum* (▲)
in N. America.